

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



109

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> C12N 15/24, C07K 14/54, 7/06, 16/34, A61K 38/20, 39/395	<b>A1</b>	<b>(11) International Publication Number:</b> WO 96/01318 <b>(43) International Publication Date:</b> 18 January 1996 (18.01.96)
<b>(21) International Application Number:</b> PCT/DK95/00227 <b>(22) International Filing Date:</b> 7 June 1995 (07.06.95) <b>(30) Priority Data:</b> 0800/94 5 July 1994 (05.07.94) DK <b>(71) Applicant (for all designated States except US):</b> NYCOMED DAK A/S [DK/DK]; Langebjerg 1, P.O. Box 88, DK-4000 Roskilde (DK). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> GRØNHØJ LARSEN, Christian [DK/DK]; Skejbygårdsvej 300, DK-8240 Risskov (DK). GESSER, Borbala [SE/DK]; Pilegårdsvej 233, DK- 8361 Hasselager, Kolt (DK). <b>(74) Agent:</b> PLOUGMANN & VINGTOFT A/S; Sankt Annæ Plads 11, P.O. Box 3007, DK-1021 Copenhagen K (DK).	<b>(81) Designated States:</b> AM, AT (Utility model), AU, BB, BG, BR, BY, CA, CH, CN, CZ, CZ (Utility model), DE (Utility model), DK (Utility model), EE, ES (Utility model), FI, FI (Utility model), GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), TJ, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).  <b>Published</b> <i>With international search report.</i> <i>With amended claims.</i>	

**(54) Title:** IMMUNOMODULATORS

**(57) Abstract**

This invention relates to a polypeptide other than human interleukin (10) which has at least one of the following properties: a) induces inhibition of spontaneous IL-8 production by human monocytes, b) induces inhibition of IL-1 $\beta$  induced IL-8 production by human peripheral blood mononuclear cells (PBMC), c) induces production of interleukin-1 receptor antagonistic protein (IRAP) by human monocytes, d) induces chemotactic migration of CD8+ human T lymphocytes *in vitro*, e) desensitizes human CD8+ T cells resulting in an unresponsiveness towards rhIL-10, f) suppresses the chemotactic response of CD4+ T human lymphocytes towards IL-8, g) suppresses the chemotactic response of human monocytes towards MCAF/MCP-1, h) does not inhibit class II MHC molecule expression on human monocytes, in contrast to human IL-10, i) induces the production of IL-4 by cultured normal human CD4+ T cells, j) reduces the TNF $\alpha$  production in human mixed leukocyte reaction. In particular, the invention relates to the nonapeptide Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn and analogues and variants thereof.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## FIELD OF INVENTION

## BACKGROUND OF THE INVENTION

BACKGROUND OF THE INVENTION

Research from the last two decades has shown the initiation, regulation and ending of inflammation within well as the regulation of growth and differentiation by a special group of signal polypeptides genes called cytokines. Cytokines are polypeptides which

cleated cells and which transmit regulatory signals between cells, thus forming a communication network between identical or different cell types of the organism. The cytokines are extremely potent mediators and active at concentrations down to  $10^{-15}$  M. Cytokines are also key factors for the development of cellular immune reactions, which in turn form the basis for the clinical manifestations of inflammation due to infection, allergy, trauma, graft vs. host reactions and auto-immune diseases. The allergic and auto-immune diseases are explained by abnormalities in the immune system, especially in the T lymphocyte-mediated immunity, but generally these diseases are of unknown etiology. *In vitro* studies, animal experiments and clinical experimental studies have shown that cytokines play important pathophysiological roles for the inflammatory reactions related to auto-immune diseases, allergy, ischemia, reperfusion injury, trauma, infections, and are important for the development of cancer, atherosclerosis, pregnancy and fetal development, bone homeostasis. Cytokines may be involved in other immunoinflammatory and proliferative diseases as will be described in further detail in the following.

The said diseases are usually chronic and the treatment is palliative, i.e. most of the drugs prescribed in connection with the said diseases are directed at the allaying of symptoms and usually have no curative effect. Other treatments are so-called substitution therapies which involve life-long supplying to the patient of substances, e.g. hormones, needed due to a reduced/insufficient internal production of said substance. Said treatments are often unsatisfactory, imply unwanted and often serious side-effects, and merely delay rather than prevent the progression of the disease. Thus, improved methods of treatments and improved pharmaceutical compositions are highly needed.

Interleukin-10 (IL-10) is a recently described natural endogenous immunosuppressive cytokine, identified in both the murine and human organism. Murine interleukin 10 (mIL-10) was

originally described as a cytokine synthesis inhibitory factor released from TH<sub>2</sub> helper T-cell clones, but it also carries proliferative effects upon various subsets of lymphocytes, including an enhancing effect upon cloning efficacy of CD4<sup>-</sup>, 8<sup>+</sup> murine splenic T cells (4). Human interleukin 10 (hIL-10) has recently been sequenced and revealed to have high homology with mIL-10 at DNA sequence level as well as on amino acid level. Furthermore, swine interleukin 10 has recently been sequenced and revealed to have high homology with human IL-10 at DNA sequence level as well as on amino acid level (88), see also Figure 2. Also, hIL-10 has high homology with an open reading frame in the Epstein-Barr virus genome, BCRF1, and viral IL-10 does show some activity similar to hIL-10, cf. Fig. 1 (5).

Human IL-10 is produced by activated T cell clones and immortalized B cells, and in addition to its cytokine synthesis inhibitory factor (CSIF) activity, inhibiting the production of several pro-inflammatory cytokines and colony-stimulating factors, it also induces the production of a natural interleukin-1 receptor antagonist protein/peptide (IRAP) by mononuclear cells, thereby indirectly inhibiting IL-1 activity. IL-10 also downregulates its own production by monocytes and inhibits the expression of class II MHC expression (12). Further, hIL-10 reduces antigen-specific proliferation of human T cells and CD4<sup>+</sup> T cell clones, when using monocytes as antigen-presenting cells. *In vivo* experiments in mice indicate that the outcome of Leishmania infection is dependent upon the cytokine profile from responding CD4<sup>+</sup> T lymphocytes (13). In C57BL/6 mice resistant to Leishmania infection, CD4<sup>+</sup> T cells from draining lymph nodes show up-regulation of IFN- $\gamma$  and IL-2 cytokines, whereas the sensitive BALB/c mice in their draining lymph nodes have CD4<sup>+</sup> responding T cells releasing IL-4 and IL-10, which could be demonstrated to correlate with disease progression (13). Thus, IL-10 may exert potent regulatory effects on immunological responses both *in vitro* and *in vivo*. Further IL-10 strongly affects chemokine biology since human interleukin 10 is a specific chemotactic

factor towards CD8+ T cells, while IL-10 suppresses the ability of CD4+, but not CD8+, T cells to migrate in response to the T cell chemotactic cytokine, IL-8 (14). IL-10 also inhibits the chemotactic effect of other chemokines MCP-1/MCAF and RANTES (75). Since IL-10 is a deactivator of monocyte/macrophage functions and an inhibitor of Th1 activity, drugs with full or partial IL-10-like activity may possess therapeutic effect in diseases characterized by imbalance in cytokine production and/or activities.

It has previously been proposed to prepare pharmaceutical compositions comprising hIL-10 or vIL-10, and the use of hIL-10 or vIL-10 for the manufacture of a pharmaceutical composition for the treatment of various conditions such as septic or toxic shock, rheumatoid arthritis, graft-vs-host disease, tissue rejection, diabetes mellitus, autoimmune disorders, leukaemia and cancer has been disclosed in e.g. WO93/02693 and WO94/04180. Moreover, IL-10 antagonists, e.g. antibodies specifically binding to IL-10, have been disclosed in e.g. EP 405 980 and WO94/06473, and it has been contemplated that such antibodies could be useful in the treatment of HIV infected patients.

#### SUMMARY OF THE INVENTION

According to the present invention, it has been found that a substance other than human interleukin 10 which has one or more of the following properties:

- a) induces inhibition of spontaneous IL-8 production by human monocytes,
- b) induces inhibition of IL-1 $\beta$  induced IL-8 production by human peripheral blood mononuclear cells (PBMC),
- 30 c) induces production of interleukin-1 receptor antagonistic protein (IRAP) by human monocytes,
- d) induces chemotactic migration of CD8+ human T lymphocytes *in vitro*,

- e) desensitizes human CD8+ T cells resulting in an unresponsiveness towards rhIL-10,
- f) suppresses the chemotactic response of CD4+ human T lymphocytes towards IL-8,
- 5 g) suppresses the chemotactic response of human monocytes towards MCAF/MCP-1,
- h) does not inhibit class II MHC molecule expression on human monocytes, in contrast to human IL-10,
- i) induces the production of IL-4 by cultured normal human  
10 CD4+ T cells,
- j) reduces the TNF $\alpha$  production in human mixed leukocyte reaction,

such as a polypeptide which comprises the amino acid sequence Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn or an analogue or variant  
15 of said sequence (a nona-peptide with sequence homology to hIL-10, called IT9302), and derivatives thereof can be used for the prevention and/or treatment of certain forms of inflammatory processes, especially forms related to the immune and/or hormone system. It is contemplated (as described in detail in the following description of immunological  
20 mechanisms) that the action mechanism is via interference with the action of mediators of the immune system, in particular cytokines such as monokines, lymphokines, chemokines and monokine-receptor antagonists, i.e. that the substance of the invention interferes with/suppresses the production  
25 and/or action of certain cytokines and thus inhibits pathological processes leading to tissue damage, and that the substance of the invention induces the production of natural monokine-receptor antagonists thus interfering with/suppressing the action of certain cytokines and thereby inhibiting  
30 pathological processes which lead to tissue damage.

An important embodiment of the present invention thus relates to a pharmaceutical composition comprising, as the active ingredient, a substance of the invention. Other embodiments  
35 of the invention are a substance which is capable of neutralizing one or more of the activities a) to g) mentioned

above, e.g. an antibody, and a pharmaceutical composition comprising such a substance.

- In a further aspect, the present invention relates to the use of a substance of the invention for the manufacture of a pharmaceutical composition for substantially inhibiting a biological effect related to a cytokine, i.e. the use of a substance of the invention as an IL-1 receptor antagonist protein/peptide, lymphokine, monokine, interleukin, interferon, chemokine or colony-stimulating factor. Another aspect relates to the use of a substance of the invention for the manufacture of a pharmaceutical composition for the prophylaxis or treatment of a condition related to the disturbance of a cytokine system, i.e. the IL-1 receptor antagonist protein/peptide, lymphokine, monokine, interleukin, interferon, chemokine or colony-stimulating factor system. In another aspect, the invention also relates to a method of treating a condition in a human related to a disturbance in a cytokine system which method comprises administering to the subject an effective amount of a substance of the invention.
- 20 The cellular immune system takes part in the development of such disorders as infectious, inflammatory and neoplastic diseases. Immunocompetent cells and their products may play important roles in the initiation, progression and possible chronic nature of development of inflammatory conditions.
- 25 These disorders are often without known etiology and includes common diseases such as diabetes mellitus, rheumatoid arthritis, inflammatory diseases of the gastro-intestinal tract and of the skin. Apart from these examples, cell-mediated immunity or pro-inflammatory mediators, however, contribute to many
- 30 other inflammatory and proliferative diseases (see Table 2).



TABLE 2

Some diseases where macrophages/T-lymphocyte-mediated immune reactions are considered pathogenetically important

---

- 5 *Skin diseases:*
  - Psoriasis
  - Atopic dermatitis
  - Contact dermatitis
  - Cutaneous T cell lymphoma (CTCL)
- 10 Sezary syndrome
  - Pemphigus vulgaris
  - Bullous pemphigoid
  - Erythema nodosum
  - Scleroderma
- 15 *Auto-immune (including rheumatic) diseases:*
  - Uveitis
  - Bechet's disease
  - Sarcoidosis Boeck
  - Sjögren's syndrome
- 20 Rheumatoid arthritis
  - Juvenile arthritis
  - Reiter's syndrome
  - Gout
  - Osteoarthritis
- 25 Systemic lupus erythematosus
  - Polymyositis
  - Myocarditis
  - Primary biliary cirrhosis
  - Crohn's disease
- 30 Ulcerative colitis
  - Multiple sclerosis and other demyelinating diseases
  - Aplastic anaemia
  - Idiopathic thrombocytopenic purpura
  - Multiple myeloma and B cell lymphoma
- 35 Simmons' panhypopituitarism
  - Graves' disease and Graves' ophthalmopathy

Subacute thyreoditis and Hashimoto's disease  
Addison's disease  
Insulin-dependent diabetes mellitus (type 1)

*Other diseases*

- 5 Various clinical syndromes with vasculitis (e.g. polyarteritis nodosa, Wegener's granulomatosis, Giant cell arteritis Fever, malaise  
Anorexia (e.g. in acute and chronic inflammatory and infectious diseases)
  - 10 Disseminated intravascular coagulation (DIC)  
Arteriosclerosis (atherosclerosis)  
Shock (e.g. in gram-negative sepsis)  
Cachexia (e.g. in cancer, chronic infectious and chronic inflammatory diseases)
  - 15 Transplant rejection and graft vs. host disease
- 

*Cytokines:*

- T-lymphocytes orchestrate the induction and regulation of cell-mediated immune reactions, and cytokine products (lymphokines) of the T cells initiate and control the immune response (1,2). The lymphocyte-activating mediators (lymphokines) produced by antigen-presenting cells belong to a group of polypeptides called cytokines. Cytokines are transmitters of cell-to-cell communication in both physiological and pathophysiological conditions, and may also function as hormones providing signals between the immune system and other tissues and organs. Cytokines may also be produced by cells outside the immune system and it is generally believed that all nucleated cells are capable of producing one or several cytokines. Thus, e.g. keratinocytes and fibroblasts are potent producers of cytokines and in this system cytokines may function as autocrine or paracrine hormones independent of the immune system (3).

*Interleukin-10:*

Murine interleukin 10 (mIL-10) was originally described as a cytokine synthesis inhibitory factor (CSIF) released from TH<sub>2</sub> helper T-cell clones but it has also proliferative effects upon various subsets of lymphocytes, including an enhancing effect upon cloning efficacy of CD4<sup>+</sup>, 8<sup>+</sup> murine splenic T cells (4). Human interleukin 10 (hIL-10) has recently been described (5) and has high homology with an open reading frame in the Epstein-Barr virus genome, BCRF1, and viral IL-10 does show some activity similar to hIL-10. In the following, the biochemical, biological, physiological and possible pathophysiological roles of IL-10 will be summarized.

#### *IL-10 structure:*

The primary structures of mouse (mIL-10) and human IL-10 (hIL-10) revealed a high degree of nucleotide sequence homology (>80%) throughout their entire length (4, 5). The only significant difference is the insertion of a human Alu repetitive sequence element in the 3'-untranslated region of the hIL-10 cDNA clone. The mIL-10 and hIL-10 cDNAs encode very similar open reading frames (ORF) of 178 amino acids, including hydrophobic leader sequences and corresponding to 73% amino acid homology. mIL-10, which is active on murine cells, does not cross-react significantly on human cells. hIL-10 is an 18 kDa polypeptide which lacks detectable carbohydrate, but mIL-10 is N-glycosylated at a site near its N-terminus which is missing from hIL-10. Both mIL-10 and recombinant hIL-10 (rhIL-10) are expressed as non-covalent homodimers. The extent to which mIL-10 or hIL-10 monomers are biologically active is not yet certain. mIL-10 and hIL-10 with polypeptide "tags" of at least 8 amino acids at the N-terminus and 21 amino acids at the C-terminus showed no detectable loss of activity according to a publication from Moore et al. who were the first to sequence human IL-10 (6). Tagging C and N terminal ends of the whole IL-10 does not necessarily cause functional changes, since tagging is occasional or incidental. The high number of possible amino acid substitutions as described in further detail in the following might also

explain while tagging failed to show missing functions. Recombinant mIL-10 and hIL-10 have been expressed in: CDS7 cells, mouse myeloma cells, chinese hamster ovary cells, a baculovirus expression system, and *E. coli*. The biological activities of these rIL-10 proteins are so far indistinguishable (6).

The mIL-10 gene comprises five exons arrayed over approximately 5.1 kb of DNA. The genomic clone itself encodes an expressible mIL-10 protein. The mIL-10 and hIL-10 genes are on mouse and human chromosome 1 (6, 7).

mIL-10 and hIL-10 exhibit strong DNA and amino acid sequence homology to an open reading frame in the Epstein-Barr virus genome, BCRF1, and the homology is confined to the mature protein coding sequence and is not detected in the signal sequence or 5'- and 3'-untranslated sequences (5). Of the three sequences, mIL-10 and hIL-10 are the more closely related pair at the DNA sequence level (81%), while the DNA sequences encoding the mature hIL-10 and BCRF1 proteins have 71% homology. The homology between hIL-10 and BCRF1 is 84% on the amino acid level. It is hypothesized that the mIL-10 and hIL-10 genes are evolved from a common ancestor, while BCRF1 represents an ancestral processed, captured cellular cytokine gene and that the BCRF1 protein has been constrained to resemble hIL-10. BCRF1 is expressed during the lytic cycle of the EBV. The BCRF1 ORF encodes a 17 kDa secreted polypeptide which, like hIL-10, contains little or no glycosylation. BCRF1 displays some activities of IL-10 and has been called viral IL-10 (vIL-10), although its activity has been found to be 10% of hIL-10.

### IL-10 activity on cytokine production:

hIL-10 inhibits the production of a number of cytokines including interferon- $\gamma$  (IFN- $\gamma$ ), Tumor Necrosis Factor- $\alpha$  (TNF- $\alpha$ ), Granulocyte Macrophage Colony Stimulating Factor (GM-CSF), Granulocyte-CSF (G-CSF), IL-1 $\alpha$ , IL-1 $\beta$ , IL-2, IL-6,

IL-8 and Monocyte Chemotactic polypeptide-1 (MCP-1/MCAF) by monocytes/macrophages and/or T-lymphocytes (4, 5). IL-10 also inhibits the ability of monocytes to migrate as a response to the chemokine MCP-1/MCAF (75). Further, hIL-10 induces the production of an endogenous, natural interleukin-1 receptor antagonist (IRAP) (6), which inhibits IL-1 $\alpha$  and IL-1 $\beta$  by competing with receptor binding. Since IL-8 is strongly inducible by IL-1 $\alpha$  and by IL-1 $\beta$ , IL-10 exerts part of its inhibitory effect on IL-8 production by stimulating the production of the IL-1-receptor antagonist IRAP. This last mechanism is of considerable importance for the present invention as described and exemplified in the following. IRAP has anti-inflammatory activities (9), and its therapeutic effect in rheumatoid arthritis has been suggested (10). Also, IRAP proved to be effective in the treatment of sepsis syndrome and a dose-dependent, 28-day survival benefit was associated with IRAP treatment ( $p = 0.015$ ) in a study by Fisher et al. (11). IRAP may exert parts of its anti-inflammatory effects by inhibiting chemokine-production such as the production of IL-8.

#### *IL-10 and antigen expression:*

IL-10 inhibits the expression of class II MHC expression on human monocytes (8). Constitutive and IL-4 or IFN- $\gamma$  induced expression of HLA-DR/DP and DQ was inhibited by hIL-10 (12). In addition, monocytes pre-incubated with IL-10 are refractory to subsequent induction of class II MHC expression by IL-4 or IFN- $\gamma$ . IL-10 inhibits class II expression by human monocytes following activation by LPS (12, 76). BALB/c mice given 1 to 10 mg of IL-10 concomitantly with a lethal dose of LPS were protected from death (6).

IL-10 inhibits nitrogen intermediates and superoxide anions. IL-10 also inhibits reactive nitrogen intermediate (NO) as well as reactive oxygen intermediates (H<sub>2</sub>O<sub>2</sub>) by macrophages following activation by IFN- $\gamma$  (13).

*IL-10 and T cell activity:*

IL-10 has also modulatory effects on T cell functions/activity. Thus, hIL-10 is a potent chemotactic factor to CD8+ T lymphocytes, while hIL-10 does not show chemotaxis towards CD4+ T cells (14). Additionally, IL-10 suppresses the capacity of CD4+ T cells to respond to the chemotactic signals of the  $\beta$ -chemokine RANTES as well as the  $\alpha$ -chemokine IL-8. hIL-10 also directly inhibits the proliferation of human peripheral blood T cells and CD4+ T cell clones (14).

*10 IL-10 and B lymphocytes:*

hIL-10 co-stimulates B-lymphocyte proliferation induced by cross-linking surface Ig with immobilized anti-IgM antibody, and this effect is enhanced when B-cells are stimulated by cross-linking of their CD40 antigen with anti-CD40 antibody and mouse L cells expressing human Fc $\gamma$ RII/CD32 (15). The effect of IL-10 on proliferation and differentiation of activated human B cells suggests that this cytokine may account for much of the superior ability of T cells expressing hIL-10 to provide help for B cell responses.

*20 IL-10 as a homeostatic factor for the immune system:*

The physiological consequences of the functions of IL-10, mentioned above, are believed to be a certain degree of homeostasis on the immune system. Thus, IL-10 clearly inhibits helper T cells functions and probably stimulates T cells with suppressor functions. Therefore, like IL-4, IL-10 is believed to regulate the balance between Th1 and Th2 cytokine profiles of T cells. Especially, it is believed that IL-10 inhibits the differentiation of Th1 cells. Since Th1 cells are characterized by the production of cytokines (IFN- $\gamma$  and IL-2) which favour cell-mediated immune responses while Th2 cells produce cytokines (IL-4, IL-5 and IL-10) which favour a humoral response and suppress a cell-mediated immune response, IL-10 is likely to suppress a T cell-mediated immune reaction such

as delayed type hypersensitivity reactions, while it favours humoral responses.

As a consequence of the features of IL-10 mentioned above, the activity of IL-10 has been described as an inhibitor of macrophage and Th1 cytokine synthesis. Therefore it has been studied whether lack of IL-10 production and/or activity may play a role in diseases where an enhanced cell-mediated immunoreactivity is believed to play a role for the disease, such as in auto-immune diseases and inflammation. Anti-IL-10 antibody-treated mice show a stronger inflammatory response to monokine-induced inflammation and are significantly more susceptible to death induced by LPS-induced septic shock, a monokine-mediated inflammatory reaction (16). Also, IL-10 knock-out mice spontaneously develop inflammatory reactions of the gut similar to that of colitis ulcerosa (17). Additionally, it has been investigated whether IL-10 plays a role in different parasite, mycobacterial or viral infections, and IL-10 has been shown to play a pathophysiological role in the immune-paresis related to the infection with *Schistosoma mansoni* (18). Also a role in *Mycobacterium leprae* infections has been suggested. Recently, it has been found that AIDS patients with a poor prognosis have a higher level of IL-10 in plasma, and it has been suggested that this contributes to the immune paresis which is known from AIDS (19).

#### 25 *Therapeutic considerations:*

These *in vivo* results/data strongly suggest a homeostatic role of IL-10 in controlling cell-mediated and monokine-amplified immune inflammation and indicate the wide-ranged therapeutical applications of IL-10 or a drug with IL-10-like activity in the treatment of diseases which are characterized by a decreased/insufficient production and/or activity of IL-10. Since the substance of the present invention exemplified by IT9302 exerts IL-10-like activity by

- 1) inducing IRAP production by human monocytes,

- 2) inhibiting spontaneous IL-8 production by human monocytes,
- 3) inhibiting IL-1 $\beta$ -stimulated IL-8 production by peripheral blood mononuclear cells (PBMC),
- 5 4) stimulating chemotactic migration of CD8+, but not CD4+, human T lymphocytes,
- 5) desensitizing human CD8+ T cells towards rhIL-10-induced chemotactic migration,
- 10 6) inhibiting IL-8-mediated human CD4+ T cell chemotaxis, and
- 7) inhibiting MCP-1/MCAF-mediated human monocyte chemotaxis,
- 8) inducing the production of IL-4 by cultured normal human CD4+ T cells,
- 15 9) reducing the TNF $\alpha$  production in human mixed leucocyte reaction,

this polypeptide and analogues thereof may thus possess the same therapeutic possibilities as hIL-10. Table 3 lists some diseases where an immune-modulator like IL-10 or an immune-

20 modulator with IL-10-like activity may have therapeutic importance:

TABLE 3

25 Some diseases where an immune-modulator with IL-10-like activity, due to its induction of IRAP production and/or inhibition of cytokine-production and/or activity may have therapeutic importance (ref. 20-74)

---

Pre-term labour caused by infection or other conditions

Rheumatoid arthritis

30 Lyme's arthritis

Gout

Sepsis syndrome

Hyperthermia

Ulcerative colitis or enterocolitis

35 Osteoporosis

Cytomegalovirus

Periodontal diseases



- Glomerulonephritis  
Chronic, non-infectious inflammation of the lung (e.g. sarcoidosis and smoker's lung)  
Granuloma formation
- 5 Fibrosis of the liver  
Fibrosis of the lung  
Transplant rejection  
Graft vs. host disease  
Chronic myeloid leukaemia
- 10 Acute myeloid leukaemia  
Other neoplastic diseases  
Asthma bronchiale  
Diabetes mellitus, type I (insulin dependent)  
Arteriosclerosis/atherosclerosis
- 15 Psoriasis  
Chronic B lymphocyte leukaemia  
Common variable immunodeficiency  
Side-effects using other biological response modifiers  
Disseminated intravascular coagulation
- 20 Systemic sclerosis  
Encephalomyelitis  
Lung inflammation  
Hyper IgE syndrome  
Enterocolitis
- 25 Cancer metastasis and growth  
Adoptive immune therapy  
Acquired respiratory distress syndrome  
Sepsis  
Reperfusion syndrome
- 30 Postsurgical inflammation  
Organ transplantation  
Alopecia
- 

#### DETAILED DESCRIPTION OF THE INVENTION

- 35 *Development of an IL-10-homologous nonapeptide with IL-10-like activity:*

Partial sequences of hIL-10 having a length of 9 amino acids was chosen according to the principle that the sequences should possess high homology between vIL-10 and hIL-10, but as low homology to mIL-10 as possible. This strategy was  
5 based on the fact that vIL-10 cross-reacts partly with hIL-10, while mIL-10 does not cross-react with hIL-10 (see above). Thus, sequences of the hIL-10 responsible for certain activities in the human organism may be located at domains where there is high homology between hIL-10 and vIL-10 but  
10 low in relation to mIL-10.

The signal polypeptide of hIL-10 is supposed to consist of the first 18 amino acids. The mature protein starts from amino acid No. 19 (No. 1 in the functional protein, which is a serine), and the total protein is containing 160 amino  
15 acids. By looking at the human and viral IL-10 COOH-terminal sequences and in particular position 157 and 159, containing lysine and arginine residues, this domain was found to be favourable for being partly responsible for receptor binding. (Figs. 1 and 2).

20 After screening several candidates, obtained by chemical synthesis, for IL-10-like activity, it was found that a synthetic nonapeptide, IT9302, possessed some immuno-suppressive activities which mimic those of hIL-10 as described in further detail in the following examples. IT9302 corresponds to  
25 a nonapeptide sequence from the C-terminal end of hIL-10 with the following amino acid sequence:

NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn-COOH (SEQ ID NO:1)

IT9302 has 100% homology with a segment of hIL-10 corresponding to amino acids Nos. 152 to 160 at the C-terminal end.

30 Furthermore, this polypeptide sequence has 6 amino acids in common with the virus vIL-10 in the corresponding area, see Figs. 1 and 2, which means that 6 of 9 or 66% of the amino acids are homologous as defined herein, and 4 amino acids in common with mIL-10, i.e. 4 of 9 or 44%. This nonapeptide is

the only sequence obtained or derived from hIL-10 and examined until now which has the special properties demonstrated in the examples. Until contemplated by the present inventors, no one has pointed to this sequence as a functional domain.

5 In its broadest aspect, the present invention thus relates to a substance which exhibits hIL-10 agonist activity, i.e. a substance other than human interleukin 10 which has one or more of the following properties:

- 10 a) induces inhibition of spontaneous IL-8 production by human monocytes,
- b) induces inhibition of IL-1 $\beta$  induced IL-8 production by human peripheral blood mononuclear cells (PBMC),
- c) induces production of interleukin-1 receptor antagonistic protein (IRAP) by human monocytes,
- 15 d) induces chemotactic migration of CD8+ human T lymphocytes *in vitro*,
- e) desensitizes human CD8+ T cells resulting in an unresponsiveness towards rhIL-10,
- f) suppresses the chemotactic response of CD4+ human T
- 20 lymphocytes towards IL-8,
- g) suppresses the chemotactic response of human monocytes towards MCAF/MCP-1,
- h) does not inhibit class II MHC molecule expression on human monocytes, in contrast to human IL-10,
- 25 i) induces the production of IL-4 by cultured normal human CD4+ T cells,
- j) reduces the TNF $\alpha$  production in human mixed leukocyte reaction.

Among these activities, d) to g) are considered the most  
30 unique. One important embodiment of the invention is a substance which exhibits hIL-10 agonist activity as defined above and which comprises the amino acid sequence Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn or an analogue or variant of said sequence. This sequence, and all other polypeptide sequences  
35 in the present specification and claims, are, also when not

explicitly stated, written from the N-terminal to the C-terminal end in the conventional format.

The nonapeptide IT9302 is very potent to induce different functions and is very stable, and it is presumed that it cannot be incorrectly coupled to receptors. A nonapeptide has been chosen because generally a 9 amino acid polypeptide sequence is unique for a protein. However, it is to be noted that the 6 amino acids at the very end of hIL-10 seem to be the most important ones. Within the scope of the present invention is thus a substance or polypeptide comprising a subsequence of the amino acid sequence Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn (SEQ ID NO:1).

It is considered likely that some amino acid substitutions will not have adverse effects on the hIL-10 agonist activity as defined herein as long as the threonine, the lysine and the arginine are present and with one amino acid placed inbetween.

In its broadest aspect, the present invention thus relates to a polypeptide having the formula

Thr-X<sub>4</sub>-Lys-X<sub>5</sub>-Arg-X<sub>6</sub> (SEQ ID NO:19),

wherein

X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and  
X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.


In a further aspect, the invention relates to a polypeptide having the formula

X<sub>3</sub>-Thr-X<sub>4</sub>-Lys-X<sub>5</sub>-Arg-X<sub>6</sub> (SEQ ID NO:20),

wherein

X<sub>3</sub>, X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and  
X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.


In yet a further aspect, the invention relates to polypeptides having the formula



wherein

X<sub>2</sub> is Tyr or Phe,  
X<sub>3</sub>, X<sub>4</sub> and X<sub>5</sub> are independently selected from the group  
consisting of Met, Ile, Leu and Val; and  
X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.

Preferred embodiments of the invention are polypeptides having the formula



wherein

X<sub>1</sub> is Ala or Gly,  
X<sub>2</sub> is Tyr or Phe,  
X<sub>3</sub>, X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and  
X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.

Examples of specific polypeptides which are presumed to have hIL-10 agonist activity as defined above are as follows:

1. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Ile-Lys-Met-Arg-Asn-COOH (SEQ ID NO:2)
2. NH<sub>2</sub>-Ala-Phe-Met-Thr-Leu-Lys-Leu-Arg-Asn-COOH (SEQ ID NO:3)
- 25 3. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Val-Arg-Glu-COOH (SEQ ID NO:4)
4. NH<sub>2</sub>-Gly-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:5)
5. NH<sub>2</sub>-Ala-Phe-Met-Thr-Met-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:6)
6. NH<sub>2</sub>-Ala-Tyr-Ile-Thr-Met-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:7)
7. NH<sub>2</sub>-Ala-Tyr-Leu-Thr-Met-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:8)

8. NH<sub>2</sub>-Ala-Tyr-Val-Thr-Met-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:9)
9. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Ile-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:10)
10. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Leu-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:11)
11. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Val-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:12)
- 5 12. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asp-COOH (SEQ ID NO:13)
13. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Met-Arg-Asp-COOH (SEQ ID NO:14)
14. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Val-Arg-Asp-COOH (SEQ ID NO:15)
15. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Ile-Asp-Gln-COOH (SEQ ID NO:16)
16. NH<sub>2</sub>-Ala-Tyr-Met-Thr-Met-Lys-Ile-Asp-Glu-COOH (SEQ ID NO:17)

10 For comparison, another nonapeptide called IT9301 with 100%  
homology to a sequence of hIL-10 was synthesized. This poly-  
peptide, IT9301, corresponds to a nonapeptide sequence start-  
ing at the N-terminal end of hIL-10, (amino acids nos. 19 to  
27, i.e. the first 9 amino acids of the mature protein) and  
15 has the following amino acid sequence:

NH<sub>2</sub>-Ser-Pro-Gly-Gln-Gly-Thr-Gln-Ser-Glu-COOH (SEQ ID NO:18)

This polypeptide showed no IL-10-like activity using the  
techniques described below and was only chosen because it is  
at the other end of hIL-10 and mostly served as a control for  
20 negative results. It does not have the characteristics of the  
C-terminal sequence with regard to sequence homology to vIL-  
10, and favourable amino acids for receptor binding. IT9301  
was tested in the IL-1 $\beta$  induced peripheral blood mononuclear  
cell system for IL-8 production, but did not induce any inhi-  
25 bition at all of the IL-8 production from the cells.

In accordance with the present invention, the term "a hIL-10  
agonist substance of the invention" comprises any pharmaceu-  
tically active and acceptable compound being identical or  
structurally similar to IT9302 and exhibiting relevant biolo-  
30 gical actions similar to those of IT9302, including derivati-  
ves of IT9302, especially pharmaceutically acceptable salts,  
esters and solvates as well as conjugates of IT9302 or of the  
IT9302 derivatives including peptido-mimetics. Covalent  
coupling of IT9302 at the NH<sub>2</sub>-terminal end to appropriate

carriers, e.g. polyethylene glycol or sugar might extend the half-life of the polypeptide *in vivo*.

The following definitions are employed in the present text:

5 "Cytokine" is a general term for a proteinaceous mediator released primarily but not exclusively by a cell population of the immune system as a response to a specific stimulating agent, e.g. a specific antigen or an alloantigen; or a non-specific, polyclonal activator, e.g. an endotoxin or other cell wall components of a gram-negative bacterium.

10 "Lymphokine" is a general term for a proteinaceous mediator released by sensitized lymphocytes as a response to a stimulating agent, e.g. a specific antigen or an alloantigen; or by a lymphocyte challenged by a polyclonal activator, e.g. an endotoxin or other cell wall components of a gram-negative  
15 bacterium.

"Interleukin" is a general term for a proteinaceous mediator released primarily but not exclusively by a macrophage, T, B, or NK cell as a response to a stimulating agent, e.g. a specific antigen or an alloantigen; or by a lymphocyte chal-  
20 lenged by a polyclonal activator, e.g. an endotoxin or other cell wall components of a gram-negative bacterium.

"Monokine" is a general term for a proteinaceous mediator released primarily but not exclusively by a mononuclear phagocyte (e.g. a monocyte or a macrophage or a Kupffer cell  
25 (liver) or a Langerhans' cell (skin) as a response to any stimulating agent.

"Chemokine" is a general term for a proteinaceous chemotactic and/or leukocyte-activating mediator released primarily but not exclusively by a cell population of the immune system as  
30 a response to a specific stimulating agent, e.g. a specific antigen or an alloantigen; or a non-specific, polyclonal activator, e.g. an endotoxin or other cell wall components of

a gram-negative bacterium and belonging to a particular gene family either the chemokine- $\alpha$  gene family or the chemokine- $\beta$  gene family.

5 "Interferon" is a general term for a proteinaceous antiviral and/or monocyte-activating mediator released primarily but not exclusively by a cell population of the immune system as a response to a virus or an interferon inducer such as a polynucleotide; in particular cells of the immune system as a response to a specific stimulating agent, e.g. a specific antigen or an alloantigen; or a non-specific, polyclonal activator, e.g. an endotoxin or other cell wall components of a gram-negative bacterium.

15 "Colony-stimulating factor" is a general term for a proteinaceous, haematopoietic colony-stimulating mediator released primarily but not exclusively by a cell population of the immune system as a response to a specific stimulating agent, e.g. a specific antigen or an alloantigen; or a non-specific, polyclonal activator, e.g. an endotoxin or other cell wall components of a gram-negative bacterium.

20 "Polypeptide" as used herein means both short peptides with a length of at least two amino acid residues and at most 10 amino acid residues, oligopeptides (11-100 amino acid residues), and longer peptides (the usual interpretation of "polypeptide", i.e. more than 100 amino acid residues in length) as well as proteins (the functional entity comprising at least one peptide, oligopeptide, or polypeptide which may be chemically modified by being glycosylated, by being lipidated, or by comprising prosthetic groups). The definition of polypeptides also comprises native forms of peptides/proteins in humans as well as recombinant proteins or peptides in any type of expression vectors transforming any kind of host, and also chemically synthesized peptides.

One embodiment of the present invention relates to a polypeptide comprising a subsequence of the sequence of human IL-10



and which has a degree of homology to vIL-10 which is 66% and a degree of homology to mIL-10 which is 44%, in particular SEQ ID NO:1. By the term "homology" is meant the identity in sequences of amino acids in segments of two or more amino acids when matched with respect to identity and position of the amino acids of the polypeptides.

Homology as the term is used herein is thus a measure of similarity between two amino acid (or nucleotide) sequences. Homology is expressed as the fraction or percentage of matching amino acids (or bases) after two sequences (possibly of unequal length) have been aligned. The term alignment is used in the sense defined in (76). Roughly, two sequences are aligned by maximizing the number of matching bases (or amino acids) between the two sequences with the insertion of a minimal number of "blank" or "null" bases into either sequence to bring about the maximum overlap.

The term "homologous" is thus used here inter alia to illustrate the degree of identity between the amino acid sequence of a given polypeptide and the amino acid sequence of IT9302. The amino acid sequence to be compared with the amino acid sequence of IT9302 may be deduced from a nucleotide sequence such as a DNA or RNA sequence, e.g. obtained by hybridization as defined in the following, or may be obtained by conventional amino acid sequencing methods. The degree of homology is preferably determined on the amino acid sequence of a mature polypeptide, i.e. without taking any leader sequence into consideration. Generally, only coding regions are used when comparing nucleotide sequences in order to determine their internal homology.

Although it is contemplated that a substantial degree of homology to hIL-10 and vIL-10 is beneficial, it is not unlikely that subsequences of the subsequence of hIL-10 which show a lower degree of homology to vIL-10, say, e.g., 50%, 55% or 60%, may also show one or more beneficial hIL-10 agonist activities. Moreover, as discussed above, it is not

deemed absolutely necessary that the degree of homology to hIL-10 is 100%. Within the scope of the present invention are also polypeptides having a lower degree of homology to hIL-10, such as 75%, 80%, 85%, 90% or 95%, although 100% is preferred.

Such polypeptides may be considered analogues of the nonapeptide. By the term "an analogue or variant thereof" is thus meant a polypeptide not having exactly the amino acid sequence Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn (SEQ ID NO:1), but still having "hIL-10 agonist activity" as defined above. Generally, such polypeptides will be polypeptides which vary e.g. to a certain extent in the amino acid composition, or the post-translational modifications e.g. glycosylation or phosphorylation, as compared to IT9302 described in the examples.

The term "analogue" or "variant" is thus used in the present context to indicate a protein or polypeptide having a somewhat different, but still similar, amino acid sequence to the amino acid sequence of IT9302, allowing for minor variations that alter the amino acid sequence, e.g. deletions, site directed mutations, insertions of extra amino acids, or combinations thereof, to generate polypeptide analogues of IT9302.

Site-directed mutagenesis provides one convenient way to effect conservative amino acid substitutions, and the like, in the native protein sequence. Alternatively, analogues may be prepared by the well known methods of liquid or solid phase polypeptide synthesis utilizing the successive coupling of the individual amino acids of the polypeptide sequence, or the polypeptide can be synthesized by the coupling of individual amino acids forming fragments of the polypeptide sequence which are later coupled so as to result in the desired polypeptide.

"Conservative" as used herein means (i) that the alterations are as conformationally neutral as possible, that is, designed to produce minimal changes in the tertiary structure of the mutant polypeptides as compared to the native protein, and (ii) that the alterations are as antigenically neutral as possible, that is designed to produce minimal changes in the antigenic determinants of the mutant polypeptides as compared to the native protein. Conformational neutrality is desirable for preserving biological activity, and antigenic neutrality is desirable for avoiding the triggering of immunogenic responses in patients or animals treated with the substances of the invention. Although it is difficult to select with absolute certainty which alternatives will be conformationally and antigenically neutral, rules exist which can guide those skilled in the art to make alterations that have high probabilities of being conformationally and antigenically neutral, see e.g. (77) and (78). Some of the more important rules include (1) replacement of hydrophobic residues is less likely to produce changes in antigenicity because they are likely to be located in the protein's interior, e.g. Berzofsky (cited above) and Bowie et al. (cited above); (2) replacement of physicochemically similar, i.e. synonymous, residues is less likely to produce conformational changes because the replacing amino acid can play the same structural role as the replaced amino acid; and (3) alteration of evolutionarily conserved sequences is likely to produce deleterious conformational effects because evolutionary conservation suggests sequences may be functionally important. In addition to such basic rules for selecting mutein sequences, assays are available to confirm the biological activity and conformation of the engineered molecules. Biological assays for the substances of the invention are described more fully in the examples. Changes in conformation can be tested by at least two well known assays: the microcomplement fixation method, e.g. (79) and (80) used widely in evolutionary studies of the tertiary structures of proteins; and affinities to sets of conformation-specific monoclonal antibodies, e.g. (81).

An important embodiment of the present invention thus relates to a polypeptide in which at least one amino acid residue has been substituted with a different amino acid residue and/or in which at least one amino acid residue has been deleted or  
5 added so as to result in a polypeptide comprising an amino acid sequence being different from the amino acid sequence or a subsequence of said amino acid sequence as defined in the following, but essentially having hIL-10 agonist activity as defined above.

10 An interesting embodiment of the invention relates to a polypeptide which is an analogue and/or comprises at least part of the polypeptide of the invention amounting in total from 10 to 100 amino acids, e.g. at least 12 amino acids, at least 15 amino acids, at least 20 amino acids or at least 30 amino  
15 acids.

In a preferred embodiment of the invention, the substance or polypeptide is used in substantially pure form. To obtain this, purification of the polypeptide may be required. Examples of the procedures employed for the purification of poly-  
20 peptides are: (i) immunoprecipitation or affinity chromatography with antibodies, (ii) affinity chromatography with a suitable ligand, (iii) other chromatography procedures such as gel filtration, ion exchange or high performance liquid chromatography or derivatives of any of the above, (iv) elec-  
25 trophoretic procedures like polyacrylamide gel electrophoresis, denaturing polyacrylamide gel electrophoresis, agarose gel electrophoresis and isoelectric focusing, (v) any other specific solubilization and/or purification techniques.

Within the scope of the present invention is also a nucleotide sequence encoding a polypeptide as defined above, in  
30 particular a nucleotide sequence encoding a polypeptide comprising or being the amino acid sequence SEQ ID NO:1, e.g. a nucleotide sequence comprising the sequence GCC TAC ATG ACA ATG AAG ATA CGA AAC (SEQ ID NO:23) or a nucleotide sequence  
35 encoding a polypeptide having a subsequence of said amino

acid sequence. Moreover, a modified nucleotide sequence which differs from the nucleotide sequence above in that at least one nucleotide has been deleted, substituted or modified or at least one additional nucleotide has been inserted so as to  
5 result in a nucleotide sequence which encodes a polypeptide having hIL-10 agonist activity is within the scope of the invention.

In the present specification and claims, the term "subsequence" designates a sequence which preferably has a size of at  
10 least 18 nucleotides, more preferably at least 21 nucleotides, and most preferably at least 24 nucleotides. In a number of embodiments of the invention, the subsequence or analogue of the nucleotide sequence of the invention will comprise at least 27 nucleotides, such as at least 30 nucleotides or at  
15 least 45 nucleotides. The polypeptide encoded by the "subsequence" should conform to at least one of the criteria a)-g) above and/or the nucleotide "subsequence" should hybridize with the nucleotide sequence comprising the sequence SEQ ID NO:23 under high stringency conditions.

20 The term "highly stringent" when used in conjunction with hybridization conditions is as defined in the art that is 5-10°C under the melting point  $T_m$ , cf. Sambrook et al, 1989, pages 11.45-11.49.

The term "analogue" with regard to the DNA fragments of the  
25 invention includes a nucleotide sequence which encodes a polypeptide identical or substantially identical to the polypeptide encoded by the DNA referred to above. It is well known that the same amino acid may be encoded by various codons, the codon usage being related, *inter alia*, to the preference  
30 of the organisms in question expressing the nucleotide sequence. Thus, one or more nucleotides or codons of the DNA fragment of the invention may be exchanged by others which, when expressed, result in a polypeptide identical or substantially identical to the polypeptide encoded by the DNA frag-  
35 ment shown above.

Furthermore, the terms "analogue" and "subsequence" are intended to allow for variations in the sequence such as substitution, insertion (including introns), addition and rearrangement of one or more nucleotides, which variations do not have any substantial adverse effect on the hIL-10 agonist activity of the polypeptide encoded by the DNA fragment or a subsequence thereof. The invention thus also includes a nucleotide sequence encoding a polypeptide having a subsequence of the amino acid sequence SEQ ID NO:1.

10 The polypeptides of the invention can be produced using recombinant DNA technology. An important embodiment of the present invention relates to an expression system comprising a nucleotide sequence of the invention. Within the scope of the invention is thus also an expression system comprising a nucleotide sequence of the invention, such as a replicable  
15 expression vector which carries and is capable of mediating the expression of a nucleotide sequence as defined above.

The organism which is used for the production of the polypeptide of the invention may be a higher organism, e.g. an animal, or a lower organism, e.g. a microorganism such as *Escherichia coli*, a yeast, a protozoan, or cell derived from a multicellular organism such as a fungus, an insect cell, a plant cell, a mammalian cell or a cell line, which carries an expression system as defined above.

25 Irrespective of the type of organism used, the DNA fragment of the invention is introduced into the organism either directly or by means of a suitable vector. Alternatively, the polypeptides may be produced in the mammalian cell lines by introducing the DNA fragment or an analogue or a subsequence thereof of the invention either directly or by means of an  
30 expression vector. The polypeptides of the invention may also be produced by chemical synthesis as discussed above.

The invention furthermore relates to a plasmid vector containing a DNA sequence coding for a polypeptide of the inven-

tion or a fusion polypeptide as defined herein. In one particular important embodiment, the DNA fragment or an analogue or subsequence thereof of the invention or a fusion DNA fragment of the invention as defined herein may be carried by a replicable expression vector which is capable of replicating in a host organism or a cell line.

The vector may in particular be a plasmid, phage, cosmid, mini-chromosome or virus. In an interesting embodiment of the invention, the vector may be a vector which, when introduced in a host cell, is integrated in the host cell genome.

The polypeptide produced as described above may be subjected to post-translational or post-synthetic modifications as a result of thermal treatment, chemical treatment (formaldehyde, glutaraldehyde etc.) or enzyme treatment (peptidases, proteinases and protein modification enzymes). The polypeptide may be processed in a different way when produced in an organism as compared to its natural production environment. As an example, glycosylation is often achieved when the polypeptide is expressed by a cell of a higher organism such as yeast or preferably a mammal. Glycosylation is normally found in connection with amino acid residues Asn, Ser, Thr or hydroxylysine.

One embodiment of the invention thus relates to a method of producing a polypeptide as defined above comprising the following steps of:

- (a) inserting a nucleotide sequence as defined above in an expression vector,
- (b) transforming a suitable host organism with the vector produced in step (a),
- (c) culturing the host organism produced in step (b) under suitable conditions for expressing the polypeptide,

(d) harvesting the polypeptide, and

(e) optionally subjecting the polypeptide to post-translational modification.

It is contemplated that antagonists effect to hIL-10 or vIL-10 can be achieved inter alia by using antibodies binding specifically to the nonapeptide IT9302, and that these can be used in therapy for neutralizing high concentration of IL-10.

In another aspect, the present invention thus relates to an antibody which binds to the polypeptide of the invention.

10 The term "antibody" refers to a substance which is produced by a mammal or more precisely a cell of mammalian origin belonging to the immune system as a response to exposure to a polypeptide antigen of the invention. In the present specification and claims "an antibody" is defined as consisting  
15 essentially of the specifically binding basic unit which consists of two heavy chains and two light chains. In its broadest aspect, however, the concept of an antibody should also include e.g. a dimer or pentamer of the basic unit.

The variant domain of an antibody is composed of variable and  
20 constant sequences. The variant part of the domain is called the idiotype of the antibody. This part of the antibody is responsible for the interaction with the antigen, the antigen binding. In the present context, the term antibody is understood as the whole antibody molecule or any fragments thereof.  
25 An antibody can be fragmented during and/or after the production. It can also be made in the fragmented form to begin with and used as such or used after joining different fragments. Especially interesting fragments are binding fragments of the antibodies of the invention, e.g. Fab or  
30 Fab' fragments.

The idiotypic (antigen binding) structure of the antibody is antigenic and can thus give rise to specific antibodies



directed against the idiotypic structure. The antibodies raised against the idio~~type~~ are called the anti-idiotypic antibodies. Such antibodies may mimic the structure of the original antigen and therefore may function as the original antigen. Such antibodies may be able to substitute the original antigen for a part or all of the functions, usability and properties of the original polypeptide of the invention.

The antibodies of the invention comprise polyclonal antibodies as well as monoclonal antibodies.

10 The antibody or fragments thereof may be of a monospecific (polyclonal) kind. The monospecific antibody may be prepared by injecting a suitable animal with a substantially pure preparation of the polypeptide of the invention. This can be followed by one or more booster injections at suitable inter-  
15 vals before the first bleeding. The animals are bled about 5-7 days after each immunization. Antibodies may optionally be isolated from the serum using standard antibody purification techniques.

The animal used for the preparation of antibodies binding to  
20 the polypeptide of the invention is preferably selected from the group consisting of rabbit, monkey, sheep, goat, mouse, rat, pig, horse and guinea pigs. The cells producing the antibodies may be spleen cells or peripheral blood lymphocytes.

25 A monoclonal antibody or fragments thereof may be raised against an essential component of the polypeptide, i.e. an epitope. The monoclonal antibody may be produced using conventional techniques (Köhler and Milstein, 1975) by use of a hybridoma cell line, or by clones or subclones thereof or by  
30 cells carrying genetic information from the hybridoma cell line producing said monoclonal antibody. The monoclonal antibody may be produced by fusing cells producing said monoclonal antibody with cells of a suitable cell line, and cloning the resulting hybridoma cells producing said monoclonal anti-

body. Alternatively, the monoclonal antibody may be produced by immortalizing an unfused cell line producing said monoclonal antibody. The monoclonal antibodies are ultimately harvested from the cell growth medium. Hybridoma cells used  
5 to make monoclonal antibody may be grown *in vitro* or in the body cavity of an animal. The monoclonal antibody or fragments thereof may also be made using the recombinant DNA techniques (Huse et al. 1989).

Monoclonal antibodies may also be made by immunizing the  
10 suitable animals with a unpurified preparation of the polypeptide of the invention. The resulting hybridoma clones secreting monoclonal antibodies should be screened for their ability to binding to polypeptide(s) or its analogue.

For purposes not requiring a high specificity, the antibody  
15 may be a polyclonal antibody. Polyclonal antibodies may be obtained, e.g. as described in Harboe and Ingild, see above. More specifically, when polyclonal antibodies are to be obtained, the polypeptide of the invention or an analogue thereof is, preferably after addition of a suitable adjuvant,  
20 such as Freund's incomplete or complete adjuvant, injected into an animal. The animals are bled regularly, for instance at weekly intervals, and the blood obtained is separated into an antibody containing serum fraction, and optionally said fraction is subjected to further conventional procedures for  
25 antibody purification, and/or procedures involving use of the purified polypeptide or an analogue thereof.

The antibody may also be an anti-anti-idiotypic antibody directed against an anti-idiotypic antibody which is an antibody directed against the site of an antibody which is reac-  
30 tive with the epitope on the antigen. The anti-idiotypic antibody may be prepared by a similar method to that outlined above for the monoclonal or polyclonal antibody.

Within the scope of the invention is an antibody which binds to a substance or polypeptide as defined above, in particular

an antibody which binds specifically to a polypeptide having the amino acid sequence Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn.

In a broad aspect, the invention thus relates to a substance which is capable of neutralizing one or more of the hIL-10 activities a) to j), i.e. having hIL-10 antagonist activity, such as an antibody having these properties. A monoclonal antibody 19F1 (82) which is capable of specifically binding to IL-10 is known and can block endogenous IL-10 produced by LPS stimulated monocytes, see Table 1 (82). This antibody recognizes the natively folded IL-10, but not necessarily a whole functional domain; however, this antibody does not specifically bind IT9302.

Within the scope of the present invention is also a pharmaceutical composition comprising a such substance as well as a pharmaceutical composition comprising a substance or polypeptide of the invention.

A very important aspect of the invention relates to a pharmaceutical composition comprising a substance having hIL-10 agonist activity or hIL-10 antagonist activity as defined above and a pharmaceutically acceptable excipient. The composition may comprise e.g. purified synthesized protein or a purified recombinant polypeptide, a monoclonal or polyclonal antibody or any other substance fulfilling the criteria a)-j) or, if having hIL-10 antagonist activity, being capable of neutralizing one or more of the hIL-10 activities a) to j).

The IL-10 agonist or antagonist used in this invention may be prepared as formulations in pharmaceutically acceptable media, for example, saline, phosphate buffered saline (PBS), Ringer's solution, dextrose/saline, Hank's solution, and glucose. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as buffering agents, tonicity adjusting agents, wetting agents, detergents, and the like. Additives may also include additional active ingredients,

e.g. bactericidal agents, or stabilizers. The amount administered to the patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the host, the manner of  
5 administration, and the like.

The pharmaceutical compositions are typically intended for transdermal or parenteral administration, e.g. intravenously, subcutaneously, or intramuscularly. Orally administrative  
10 forms are also desired and can be provided by modifying the composition to bypass the stomach environment. The composition can be used for prophylactic and/or therapeutic treatment. Preferably, the pharmaceutical compositions are administered intravenously. Thus, the invention provides compositions which comprise an IL-10 agonist or antagonist substance  
15 dissolved or suspended in an acceptable carrier, preferably an aqueous carrier. These compositions may be sterilized by conventional sterilization techniques, or may be sterile filtered.

The resulting aqueous solutions may be packaged for use as  
20 is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The IL-10 agonist or antagonist may also be administered with a second biologically active agent, such as a standard chemotherapeutic agent. Such agents include but are not limited to  
25 vincristine, daunorubicin, L-asparaginase, mitoxantrone and amsacrine.

In therapeutic applications, the pharmaceutical compositions are administered to a patient in an amount sufficient to produce the desired effect, defined as a "therapeutically  
30 effective dose". The therapeutically effective dose of a IL-10 agonist or antagonist will vary according to, for example, the particular use for which the treatment is made, the manner of administration, the health and condition of the patient, and the judgment of the prescribing physician. For  
35 example, the dose for continuous infusion will typically be

in the range of about 500 ng to about 800  $\mu$ g per day for a 70 kg patient, preferably between about 10  $\mu$ g and about 300  $\mu$ g. The dose will typically be between 700 ng/kg/day and 16  $\mu$ g/kg/day.

- 5 The concentration of IL-10 agonist or antagonist in the pharmaceutical formulations can vary widely, i.e. from about 10  $\mu$ g to about 5 mg/ml, preferably between about 100  $\mu$ g and about 2 mg/ml. The concentration will usually be selected primarily by fluid volumes, viscosities, etc., in accordance  
10 with the particular mode of administration selected. Thus, a typical pharmaceutical composition for intravenous infusion could be made up to contain 250 ml of dextrose/saline solution and 2.5 mg of IL-10 agonist or antagonist.

- For solid compositions, conventional non-toxic solid carriers  
15 may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable non-toxic composition is formed by incorporating normally employed excipients, such as those carriers  
20 previously listed, and generally 10-95% of active ingredient, that is, an IL-10 agonist or antagonist substance, preferably 25-75%.

- For aerosol administration, the IL-10 agonist or antagonist  
25 is preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of IL-10 agonist or antagonists are 0.01-20% by weight, preferably 1-10%. The surfactant must, of course, be non-toxic, and preferably soluble in the propellant. Representative of such agents are  
30 the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride such as, for example, ethylene glycol, glycerol, erythritol,  
35 arbitol, mannitol, sorbitol, the hexitol anhydrides derived

from sorbitol, and the polyoxyethylene and polyoxypropylene derivatives of these esters. Mixed esters, such as mixed or natural glycerides may be employed.

5 The surfactant may constitute 0.1-20% by weight of the composition, preferably 0.25-5%. The balance of the composition is ordinarily propellant. Liquified propellants are typically gases at ambient conditions, and are condensed under pressure. Among suitable liquified propellants are the lower alkanes containing up to 5 carbons, such as butane and propane;  
10 and preferably fluorinated or fluorochlorinated alkanes. Mixtures of the above may also be employed. In producing the aerosol, a container equipped with a suitable valve is filled with the appropriate propellant, containing the finely divided polypeptide(s) and surfactant. The ingredients are thus  
15 maintained at an elevated pressure until released by action of the valve.

To enhance the serum half-life, the IL-10 agonist or antagonist may be encapsulated, introduced into the lumen of liposomes, prepared as a colloid, or other conventional techniques  
20 may be employed which provide an extended lifetime of the polypeptides. Thus, in certain embodiments, the IL-10 agonist or antagonist may be encapsulated in a liposome. A variety of methods are available for preparing liposomes, as described in, e.g., (83), (84), (85) and (86).

25 One important embodiment of the invention thus relates to the use of a substance for diminishing or neutralizing high concentration of hIL-10 and/or vIL-10, such as the use of a substance having hIL-10 antagonistic properties for the manufacture of a pharmaceutical composition for the treatment or  
30 prophylaxis of ovary cancer and/or AIDS (87, 19).

Within the scope of the invention is also a method of treating and/or preventing ovary cancer and/or AIDS, the method comprising administering, to a patient in need thereof, a therapeutically or prophylactically effective amount of a

hIL-10 antagonist substance as well as use of a compound which has an hIL-10 antagonist activity for the manufacture of a pharmaceutical composition for treatment or prophylaxis of ovary cancer and/or AIDS.

5 *Use of the substance according to the present invention*

In accordance with the present invention, it has as described above been found that IT9302 and analogues and variants thereof are useful for preventing effects of cytokines known to be pathogenetically involved in the previously described  
10 pathological conditions.

Therefore, the potentials of therapy by using the polypeptide of the invention or analogues or derivatives thereof is contemplated and should be investigated in all diseases where a therapeutic effect of hIL-10 and/or IRAP is expected (see  
15 above, Table 3).

Very important embodiments of the invention relate to use of a substance or polypeptide of the invention for the treatment or prophylaxis of one or more of the diseases mentioned in Table 3, to the use of a substance or polypeptide according  
20 to the invention for the manufacture of a pharmaceutical composition for the treatment or prophylaxis of one or more of the diseases mentioned in Table 3, as well as to a method of treating and/or preventing one or more of the diseases mentioned in Table 3, the method comprising administering to a  
25 patient in need thereof a therapeutically or prophylactically effective amount of a substance or polypeptide according to the invention.

One important aspect of the invention is thus the use of IT9302 or a functional derivative thereof for the manufacture  
30 of a pharmaceutical composition for substantially inhibiting a biological effect in a human related to a cytokine, such as a lymphokine, interleukin, monokine, chemokine, interferon, colony-stimulating factor, prostaglandin and/or leukotriene,

for the prophylaxis or treatment of a condition related to a disturbance of a cytokine system such as the lymphokine, interleukin, monokine, chemokine, interferon or colony-stimulating factor system and/or a disturbance in the prostaglandin and/or leukotriene system. As used herein, the term  
5 "pharmaceutical composition" comprises any composition suitable for human use as described in detail above.

The invention particularly relates to the use of IT9302 or a functional derivative thereof

- 10 - for substantially *inhibiting* a biological effect in a human related to a cytokine for the prophylaxis or treatment of a condition related to a disturbance of a cytokine system; and/or
- for substantially *inhibiting* a biological effect in a human  
15 related to a lymphokine for the prophylaxis or treatment of a condition related to a disturbance of a lymphokine system; and/or
- for substantially *inhibiting* a biological effect in a human related to a interleukin for the prophylaxis or treatment  
20 of a condition related to a disturbance of a interleukin system; and/or
- for substantially *inhibiting* a biological effect in a human related to a monokine for the prophylaxis or treatment of a condition related to a disturbance of a monokine system;  
25 and/or
- for substantially *inhibiting* a biological effect in a human related to a chemokine for the prophylaxis or treatment of a condition related to a disturbance of a chemokine system;  
30 and/or
- for substantially *inhibiting* a biological effect in a human related to a lymphokine for the prophylaxis or treatment of



a condition related to a disturbance of a lymphokine system; and/or

- for substantially *inhibiting* a biological effect in a human related to an interferon for the prophylaxis or treatment  
5 of a condition related to a disturbance of an interferon system; and/or
- for substantially *inhibiting* a biological effect in a human related to a colony-stimulating factor for the prophylaxis or treatment of a condition related to a disturbance of a  
10 colony-stimulating factor system; and/or
- for substantially *inhibiting* a biological effect in a human related to a prostaglandin for the prophylaxis or treatment of a condition related to a disturbance of a prostaglandin system; and/or
- 15 - for substantially *inhibiting* a biological effect in a human related to a leukotriene for the prophylaxis or treatment of a condition related to a disturbance of a leukotriene system.

#### LEGENDS TO FIGURES

- 20 Fig. 1 shows a comparison of the predicted amino acid sequences of mIL-10, hIL-10, and BCRFI. Amino acid sequence identities are indicated by vertical lines.

Fig. 2 shows the COOH terminal polypeptide sequences of IL-10 including 9 amino acids and comparing porcine, human, BCRFI  
25 and mouse proteins.

Fig. 3 is a diagram showing that IT9302 inhibits spontaneous IL-8 production by purified cultured human monocytes.

Fig. 4 is a diagram showing that IT9302 inhibits IL-1 induced (1 ng/ml) IL-8 production by human peripheral blood mononuclear cells.

Fig. 5 illustrates IRAP production by IT9302-stimulated human monocytes.

Fig. 6 illustrates IRAP production by IL-10-stimulated human monocytes.

Fig. 7 illustrates the chemotactic activity of IT9302 on CD8+ T cells

Fig. 8 illustrates the desensitization of CD8+ T cells by IT9302, resulting in unresponsiveness of CD8+ T cells towards IL-10 (10 ng/ml) induced chemotaxis.

Fig. 9 illustrates the suppression of IL-8 activity by IT9302.

Fig. 10 is a diagram showing that IT9302 inhibits MCAF/MCP-1 induced monocyte chemotaxis.

Fig. 11 shows IL-4 production in CD4+ T cell cytosolic fractions by ECL - Western Blotting.

Fig. 12 shows TNF- $\alpha$  production in human mixed lymphocyte culture cytosolic fractions by ECL - Western Blotting. TNF- $\alpha$  Western Blotting was carried out as the IL-4 described in Materials and Methods, but using a rabbit anti-human TNF- $\alpha$  antibody (Pepro Tech. Inc., London, England) and a horseradish peroxidase-labelled secondary antibody (Cat.no. P 217, Dako, Denmark).

Fig. 13 shows that LPS induced shock and leukopenia are modulated by IT9302, shown by total leukocyte counting.

## EXAMPLES

**Materials and Methods***Cytokines and chemoattractants*

Recombinant hIL-10 was obtained from Pepro Tech Inc., NJ.  
5 (Cat.No. 200 10). Recombinant hIL-1 $\beta$  and recombinant hIL-8  
were a kind gift from Dainippon Pharmaceutical Company,  
Osaka, Japan. The culture medium was RPMI 1640 GIBCO, LPS-  
free according to the Limulus Amoebocyte Lysate assay (Sigma  
E-TOXATE Kit Cat.No. 210-A1). rhMCAF/MCP-1 was a kind gift  
10 from professor Kouji Matsushima, Kanazawa, Japan.

*Leukocyte chemotaxis assay**T cell chemotaxis*

CD4+ and CD8+ T lymphocyte subsets characterized by expres-  
sing either CD4 or CD8 antigens were purified from hepari-  
15 nized blood of normal donors. Thus, peripheral blood mononu-  
clear cells (PBMC) were purified from the heparinized blood  
by diluting 100 ml of the blood with Hanks balanced salt  
solution (HBSS) 1:1 and then separated by layering the cells  
on Lymphopac™ (Nycomed Pharma, Oslo, Norway) followed by gra-  
20 dient centrifugation at 2000 rpm for 20 minutes. The mononu-  
clear cells were washed 3 times in HBSS and the cell pellet  
was diluted in 4 ml of HBSS containing 1% fetal calf serum  
and sorted at 4°C by using Dynabeads coated with monoclonal  
antibody towards CD4 or CD8 antigen (Dynabeads M-450 CD4  
25 Cat.No. 111.16, Dynabeads M-450 CD8 Cat.No. 111.08,  
DETACHaBEAD Cat.No. 125.04). The bead:cell ratio was 10:1 and  
the incubation time 1 hour. The beads were detached by adding  
polyclonal anti-mouse antibody according to the manufactu-  
rer's instructions.

30 The chemotaxis assay was a 48-well micro chamber technique  
(Neuroprobe, Rockville, MD) as previously described (74; see

ref. 3 and ref. 14). Chemoattractants were diluted in RPMI 1640 (GIBCO Cat.No. 61870-010) with 1% sterile filtered fetal calf serum and placed in the lower 25  $\mu$ l chamber. In the case of determining T cell chemotaxis, T cells ( $5 \times 10^6$ /ml) were  
5 suspended in medium and 50  $\mu$ l was placed in the upper chamber separated from the lower chamber by a 5  $\mu$ m pore-size polycarbonate, polyvinylpyrrolidone-free filter (Nucleopore Corp., Pleasanton, CA) coated with type IV collagen (Sigma Cat.No. C 0543). Cells were allowed to migrate for 2 hours at  
10 37°C at 5% CO<sub>2</sub>. The filters were then carefully removed, fixed in 70% methanol, and stained for 5 minutes in Coomassie's Brilliant Blue. Cells attached to the lower surface of the filter were counted by measuring their area using a video camera on the microscope connected to a computer system for  
15 digital analysis and supported by software for objective determination of chemotactic migration. Approximately 5% of the T cells will migrate spontaneously corresponding to between 12,000 and 13,000 cells; this may vary from day to day, but very little in the same day's experiments. As has  
20 been described earlier (ref. 3 and ref. 14), it was therefore chosen to report the results as a ratio between number of cells migrating in the sample and in the negative control, which reflects spontaneous migration. This ratio is referred to as the chemotactic index (CI). All samples were analyzed  
25 in triplicates and cell migration in each well was measured in three fields before the median value of area was estimated. In some experiments the chemotaxis membrane was not coated with collagen, and in the present assay system migrating cells will therefore drop to the bottom of the lower  
30 well of the chemotaxis chamber.

In one experiment, the chemotactic activity of IT9302 on CD8+ T cells was performed by testing serial dilutions of IT9302 added to the lower chamber and evaluating chemotaxis as described above.

35 In a second experiment, the ability of IT9302 to desensitize the migration of CD8+ T cells as a response to rHIL-10

(10 ng/ml) was studied by adding IT9302 to the target cells 30 minutes before chemotaxis. IT9302 was added in serial concentrations and the chemotactic response of rhIL-10 was evaluated as described above.

- 5 In a third experiment, the ability of IT9302 to suppress the chemotactic response of CD4<sup>+</sup> T cells towards rhIL-8 (10 ng/ml) was studied by adding IT9302 to the target cells 30 minutes before performing chemotaxis. IT9302 was added in serial concentrations and the chemotactic response of rhIL-8  
10 was evaluated as described above.

#### *Monocyte chemotaxis*

- Monocyte chemotaxis was measured using the same Boyden chamber equipment as described for T cells above. The chemo-attractant MCAF/MCP-1 was diluted in RPMI 1640 medium with  
15 0.5% BSA and added to the lower chamber at a concentration of 10 ng/ml. Monocytes, purified by the standard plastic adherence technique, from normal human PBMC, obtained as described above were suspended in RPMI 1640 medium with 0.5% BSA and then incubated for 30 minutes in the presence of IT9302  
20 at different concentrations. Subsequently, the cells were added to the upper chemotaxis chambers at a concentration of 10<sup>6</sup> cells/ml. The upper and lower chambers were separated by an 8 µm pore size polyvinyl pyrrolidone-free polycarbonate filter (Nucleopore, Pleasanton, CA). The chamber was incubated  
25 at 37°C for 90 minutes. The membranes containing migrating cells were treated as described above and a chemotactic index calculated according to the technique described above.

#### *Production of IL-8 by normal human peripheral mononuclear cells (PBMC)*

- 30 PBMC was purified from heparinized blood of normal human donors. Following gradient centrifugation with Lymphoprep™ (Nycomed Pharma, Oslo, Norway), the mononuclear cells were

diluted to  $2 \times 10^6$  cells/ml in LPS-free RPMI 1640 medium (Gibco Cat.No. 6187-010) containing 1% sterile filtered heat-inactivated fetal calf serum and penicillin (10,000 IE/ml), streptomycin (10 mg/ml) and gentamycin (2.5 mg/ml). Cells were cultured in 24 wells Nunc Micro Plates (Nunc, Denmark) and in the presence of different concentration of IT9302 (0, 1  $\mu$ g, 100 ng, 10 ng, 1 ng, 0.1 ng, 0.01 ng/ml) for 24 hours. Following 24 hours of incubation, another dose of IT9302 was added once more, and 1 hour later r-hIL-1 $\beta$  (1 ng/ml) was added to the cell cultures. Supernatants were collected after a total of 48 hours of incubation, and the concentration of the secreted IL-8 was measured by IL-8 ELISA by using an IL-8 ELISA Kit (Dainippon Pharmaceutical Co. Ltd, Osaka, Japan). Briefly, standards and cell supernatants were incubated for one hour and in duplicates at 20°C on a micro-plate shaker. Then after washing, a second antibody was added for one hour, followed by one hour of incubation with peroxidase-labelled goat anti-rabbit IgG. After washing, the reaction was developed with O-phenylenediamine. Thirty minutes later the reaction was stopped with 1.6 N sulphuric acid. Optical density (OD) was measured in an ELISA reader at 490 nm. IL-8 concentration was calculated by a calibration curve of absorbance of unknown vs concentrations of IL-8 standards.

#### *Determination of IRAP concentration*

PBMC was purified as described above. PBMC was cultured in RPMI 1640, 10% sterile heat-inactivated fetal calf serum (including penicillin 10,000 IE/ml, streptomycin 10 mg/ml, gentamycin 2.5 mg/ml) and the cell concentration was  $5 \times 10^6$  cells/ml. The monocytes were then purified by standard plastic adherence technique. Monocytes were then cultured in RPMI 1640 with 2% FCS ( $2.5 \times 10^6$  cells/ml) and with different dilutions of rhIL-10 or IT9302. The cells were stimulated for 24 hours and the supernatants were collected for IRAP determination. IRAP ELISA was carried out by using Human IL-1ra Quantikin Immunoassay Kit from R&D Systems Europe Ltd. (Cat.No. DRA 00, Abingdon, Oxon, UK).

*Determination of class II MHC antigen expression on monocytes*

PBMC was purified as described above and monocytes isolated by plastic adherence technique. Monocytes were then cultured in RPMI 1640 with 2% FCS including penicillin 10,000 IE/ml, streptomycin 10 mg/ml, gentamycin 2.5 mg/ml at a cell concentration of  $3 \times 10^6$ /ml. Cells were stimulated for 40 hours in microwells (Nunc, Denmark) with 100 ng/ml rhIL-10, or IT9302 1  $\mu$ g/ml, 100 ng/ml, 10 ng/ml. At the end of stimulation, the supernatants were removed and the cells were detached from the plastic by freezing at  $-20^\circ\text{C}$  for 20 minutes. The cells were collected in 1 ml of cold HBSS with 1% FCS and sorted at  $20^\circ\text{C}$  by using 50  $\mu$ l/ml Dynabeads M 450 coated with monoclonal antibody for HLA class II  $\beta$  chain (Cat. No. 210.03). After 20 minutes of incubation, the cells were washed 3 times with cold HBSS with 1% FCS and collected on a magnetic separation device. The cells were diluted in the above buffer and stained with methylene blue, and the rosetting cells (cells carrying Dynabeads with HLA class II mAb) were counted.

*Determination of IL-4 production by CD4+ T lymphocytes**20 Cell cultures*

CD4+ T lymphocytes were purified from heparinized normal human blood. Following Lymphoprep™ (Nycomed Pharma, Oslo, Norway) gradient centrifugation, the mononuclear cells were further sorted at  $4^\circ\text{C}$  using Dynabeads (Dynal AS, Norway) coated with monoclonal anti-CD4 antibodies. The beads were detached by adding polyclonal anti-mouse antibody (Dynal AS, Norway). The purity of the positively selected cells were higher than 99% as judged by FACS analysis. When examining the *de novo* production of IL-4 by IL-8-stimulated T cells, T cells were cultures,  $5 \times 10^6$  cells/ml in LPS-free RPMI 1640 (Gibco Cat.no. 61870-010) containing 1% sterile-filtered, heat-inactivated fetal calf serum (FCS), penicillin (10,000 IU/ml), streptomycin (10 mg/ml) and gentamycin (2.5 mg/ml).

T cells were stimulated for 3 days using rIL-8 (100 ng/ml), rIL-10 (100 ng/ml), IT9302 (10 ng/ml) and IFN- $\gamma$  (10 ng/ml). Recombinant human IL-8 (rh IL-8) was a kind gift from Dainippon Pharmaceuticals Co. Ltd., Osaka, Japan), and IFN- $\gamma$  was  
5 purchased from Boehringer Ingelheim Am Rhein, Germany. To obtain specific inhibition of IL-8 stimulation, a neutralizing monoclonal anti-IL-8 antibody (WS.4) was used (a kind gift from Dr. K. Matsushima, Japan). Recombinant IL-10 was purchased from Pepro Tech. Inc. (London, England).

10 *Preparation of cell material and culture supernatant for gel electrophoresis*

Cultured T cells and culture media were separated by centrifugation at 2000 rpm for 5 minutes. The supernatants were freeze-dried and then dissolved in 100  $\mu$ l of lysis buffer.  
15 The cells were resuspended directly in 100  $\mu$ l of gel lysis buffer (9). The material was kept at -80°C until further examination.

*ECL-Western blotting of CD4+ T cell derived proteins*

Cells or freeze-dried cell culture supernatants were used for  
20 IL-4 protein content determination. Proteins from one-dimensional 15% SDS-PAGE gels were transferred by blotting onto Hybond-ECL nitrocellulose membranes (Amersham RPN 2020D, UK) and blocked with 5% bovine serum albumin (Sigma) in Tris buffer saline pH 7.8 containing 0.1% Tween-20. The blots were  
25 then incubated with a polyclonal goat anti-human IL-4 antibody (R&D Systems, UK) followed by a horseradish peroxidase-labelled secondary antibody (Cat.no. RPN 2106 ECL, Amersham, UK), and the immunostaining was detected by exposing a film (Kodak X-OMAT-S, USA) for 90 seconds.

30 *Specificity of IT9302 amino acid sequence*

Searching for possible homology of the IT9302 sequence with other known proteins was carried out by searching on the EMBO



protein database kindly assisted by Dr. Henrik Leffers, Institute for Medical Biochemistry, University of Aarhus, Denmark.

*Amino acid sequence specificity of IT9302*

- 5 According to information from the EMBO protein database (Heidelberg) performed on 10 June 1994, IT9302 has 100% homology with a sequence of the human IL-10 and 75% with a sequence of an Epstein-Barr virus derived protein identical with vIL-10. Other virus proteins as phage T7 and tomato  
10 yellow leaf curl virus revealed 75% and 85% identity, respectively.

*Results*

- The conception of the invention took place in October and November 1992, when a nonapeptide designated IT9302 was de-  
15 signed according to the strategy outlined above, and the chemical synthesis of the first prototype (IT9302) was decided and ordered on 27 November 1992. It was expected that this nonapeptide should exert immunomodulatory activities of which some would mimic IL-10-like activity, and the following con-  
20 trol experiments (examples) were planned during this period as well. The chemical synthesis of IT9302 was carried out by using an automatic polypeptide synthesizer performed by Carlbiotech Ltd. A/S, Denmark as a payed order by the inven-  
25 tors. Afterwards the protein was purified on HPLC and the purity confirmed to exceed 95% by HPLC. This confirmed that the synthesized product was identical with the IT9302 designed by the inventors in October and November 1992.

## EXAMPLE 1

*IT9302 induced inhibition of spontaneous IL-8 production by human monocytes*

The test was performed as described in "Production of IL-8 by normal human peripheral mononuclear cells (PBMC)". Monocytes were purified by plastic adherence technique and  $3.0 \times 10^6$  cells/ml were stimulated for 40 hours. As shown in Fig. 3, IT9302 inhibited the production of IL-8 by monocytes, and at 0.1 ng/ml of IT9302 the IL-8 production was suppressed to 35% of the spontaneous production *in vitro*. The viability of cells always exceeded 80% after 1 day in culture and the addition of IT9302 did not in this or in the following examples affect viability at any concentration of IT9302 between 0.1 and 1000 ng/ml (IT9302 MW: 1,127 dalton, rhIL-10 predicted MW: 18,400 dalton).

## EXAMPLE 2

*IT9302 induced inhibition of IL-1 $\beta$  induced IL-8 production by human peripheral blood mononuclear cells (PBMC)*

The test was performed as described in "Production of IL-8 by normal human peripheral blood mononuclear cells (PBMC)". As shown in Fig. 4, IT9302, in a dose dependent manner, inhibited the IL-1 $\beta$  induced production of IL-8 by human peripheral blood mononuclear cells *in vitro*. The suppression of IL-8 production plateaued at IT9302 concentrations between 0.01 and 100 ng/ml.

## EXAMPLE 3

*IT9302 induced production of interleukin-1 receptor antagonist protein (IRAP) by human monocytes*

The test was performed as described in "Determination of IRAP concentration". As shown in Fig. 5, IT9302 dose-dependently

induced the production of IRAP by human monocytes. The production drastically increased when using concentrations of IT9302 above 10 ng/ml. Fig. 6 shows the induction of IRAP by rhIL-10 and since hIL-10 is approximately 20 times larger than IT9302, 10 ng/ml of IT9302 equals 200 ng/ml of IL-10 in molarity. Therefore the potencies of IT9302 and rhIL-10 are comparable and approximately equal with respect to the induction of IRAP at lower concentrations. At IT9302 concentrations exceeding 10 ng/ml, the induction of IRAP drastically increased and reached levels of approximately 700 ng/ml.

#### EXAMPLE 4

##### *The chemotactic effect of IT9302 on human CD8+ T lymphocytes*

The experiment was carried out as described in "Leukocyte chemotaxis assay". As shown in Fig. 7, IT9302 induced the chemotactic migration of CD8+ human T lymphocytes *in vitro*, while there was no effect on CD4+ T cells (data not shown). Again, the potency of IT9302 shown in this experiment is comparable with that of rhIL-10 shown previously (ref. 14).

#### EXAMPLE 5

*IT9302 desensitizes human CD8+ T cells resulting in an unresponsiveness towards rhIL-10*

The experiment was carried out as described in "Leukocyte chemotaxis assay". IT9302 was added to a suspension of CD8+ T cells 30 minutes before these cells were tested towards their chemotactic response to rhIL-10. As shown in Fig. 8, the pre-incubation of cells with IT9302 results in a suppressed responsiveness of the CD8+ T cells towards hrIL-10. This indicates that IT9302 may affect the binding of rhIL-10 to the IL-10 receptor.

## EXAMPLE 6

*IT9302 suppresses the chemotactic response of CD4+ T lymphocytes towards IL-8*

5 The experiment was carried out as described in "Leukocyte chemotaxis assay". As shown in Fig. 9, IT9302, in a dose-dependent manner and added to a suspension of human CD4+ T lymphocytes, inhibits the response of the CD4+ T cells towards IL-8.

## EXAMPLE 7

10 *IT9302 suppresses the chemotactic response of human monocytes towards MCAF/MCP-1*

The experiment was carried out as described in "Monocyte chemotaxis". As shown in Fig. 10, IT9302, in a dose-dependent manner and added to a suspension of human monocytes, inhibits  
15 the chemotactic response of the monocytes towards MCAF/MCP-1.

## EXAMPLE 8

*IT9302 does not inhibit class II MHC molecule expression on human monocytes*

The experiment was carried out as described in Materials and  
20 Methods and show that rhIL-10 inhibits class II MHC expression but IT9302 does not, as shown below in Table 4.

TABLE 4

Stimulation	Number of rosetting monocytes
0	$12.0 \pm 1.0 \times 10^4$
5 100 ng/ml rhIL-10	$4.6 \pm 1.4 \times 10^4$
1 $\mu$ g/ml IT9302	$11.0 \pm 3.0 \times 10^4$
100 ng/ml IT9302	$14.4 \pm 2.4 \times 10^4$
10 ng/ml IT9302	$11.2 \pm 3.2 \times 10^4$

#### 10 Discussion related to the experiments

The present data demonstrate a dose-dependent inhibitory effect of the synthetic nonapeptide, IT9302, on processes which reflect pro-inflammatory activity, including IL-8 production and monocyte and/or T cell migration. Thus, IT9302 was able to suppress the spontaneous production of IL-8 by human monocytes cultured overnight. This could be explained by a direct inhibitory effect on IL-8 mRNA production and/or subsequent protein production and or release. Another mechanism could be explained by the fact that monocytes cultured *in vitro* will express and produce IL-1, which then in turn induces the production of IL-8. This is supported by the fact that it has been demonstrated that IT9302 potently induces the production of IRAP from monocytes. IT9302 may therefore also inhibit spontaneous IL-8 production by interfering with the activity of IL-1. The observed IRAP induction by IT9302 appears to induce a biologically active IRAP, since IT9302 added to the cultures counteracts IL-1-induced IL-8 production, but only when added at least 16 hours before adding IL-1 to the cultures. This could mean that IT9302 inhibits IL-1-induced IL-8 production by inducing the production of IRAP, which then in turn blocks the activity of IL-1 through its receptor. If IT9302 directly inhibited IL-8 production, it would have been expected that addition of IT9302 to the cultures 1 hour before adding IL-1 should inhibit IL-8 production, which was not the case (data not shown). Therefore,

the observed inhibition of IL-8 production of IT9302 is likely to be due to an induction of IRAP production rather than a direct inhibition of IL-8 production. These functions of IT9302 can also be found for hIL-10 indicating that IT9302 possesses IL-10-like activities. IT9302 also mimics IL-10 activity by suppressing the ability of CD4+ T cells to migrate as a response to IL-8. Since IL-8 is related to many different states of inflammation and since CD4+ T cells appear early in the infiltrate of T cell-mediated immune inflammation such as allergy of the skin, this feature may prove to have significant therapeutic value for the control of T cell-mediated immune inflammation.

The demonstrated CD8+ T cell chemotactic activity of IT9302 is also parallel to that of IL-10, and IT9302 may thus activate T cell populations with suppressor activity contributing to the ending of T cell-mediated immune inflammation. Therefore IT9302 according to the examples which are demonstrated above, possesses therapeutic value in diseases where IL-10 and/or IRAP may also have therapeutic value. Additionally, IT9302 may have therapeutic value in diseases where IL-8 and/or MCAF and/or IL-1 are believed to have pathogenetic roles.

#### EXAMPLE 9

*IT9302 and IL-10 induce the production of IL-4 in CD4+ T lymphocytes*

#### Background

Like IL-10, IL-4 is a product of CD4+ T cells of T<sub>H</sub>2 type. It was observed that recombinant human IL-10 induces the production of IL-4 by cultured human CD4+ T cells. This means that IL-10, in addition to its own immunosuppressive functions, also induces the production of another immunosuppressive cytokine, IL-4. It was therefore tested whether IT9302 also induces the production of IL-4 by CD4+ T cells.

Thus, CD4+ T cells, purified as described in "Methods for T cell chemotaxis", and cultured as described in the section "Determination of IL-4 production by CD4+ T lymphocytes", were stimulated for 3 days with IT9302 (10 ng/ml) or IL-10 (100 ng/ml). Cytosolic fractions were collected and analyzed for their IL-4 content using Western blotting (Fig. 11) and a goat anti-human IL-4 polyclonal antibody.

As demonstrated in Fig. 11, it was observed that IL-10 as well as IT9302 induce the production of IL-4 by cultured normal human CD4+ T cells.

#### EXAMPLE 10

*IT9302 inhibits the production of TNF- $\alpha$  during a mixed leukocyte reaction (MLR)*

It has been demonstrated that the mixed leukocyte reaction is partly dependent on the production of TNF- $\alpha$  during the reaction. It has been shown that IT9302 does not significantly reduce the MLR, but it was found that there is a significant reduction in the production of TNF- $\alpha$  during the MLR.

Thus, MLR was performed by purifying human leukocytes and then culturing 1 million cells/ml from allogeneic donors for 4 days. Before establishing the cultures, one group of cells was irradiated for 2 minutes using beta irradiation. Cytosolic protein fractions were purified as described in the section "Determination of IL-4 production by CD4+ T lymphocytes", and Western blotting was performed using a rabbit anti-human TNF- $\alpha$  antibody.

As demonstrated in Fig. 12, a significant reduction in the production of TNF- $\alpha$  was observed during a human mixed leukocyte reaction.

## EXAMPLE 11

*The modulation of the LPS induced shock and leukopenia in swine using IT9302*

Since it was found that IT9302 modulates cytokine production, including TNF- $\alpha$  and IL-8, and supported by the published sequence of porcine IL-10 (Blanco et al., 1995) (see Fig. 2 for homology to the COOH-terminal peptide), it was tested whether IT9302 was able to modulate the course of LPS induced leukopenia in swine (Fig. 13).

10 In a preliminary experiment, it was tested how the intravenous injection of IT9302 0.1 mg/kg modulated the effect of intravenous injection of 2  $\mu$ g/kg LPS in swine (N = 3). IT9302 was injected 30 minutes before injection of LPS, and blood samples were drawn as described in Fig. 13. Total leukocyte count as well as differential cell count were determined, and the total number of neutrophilic leukocytes was calculated on the basis of these results.

As demonstrated, it was observed that injection of LPS caused a transient leukopenia. Injection of IT9302, however, prevented leukopenia as demonstrated in the figure.

## EXAMPLE 12

*Antibody against the synthetic polypeptide IT9302*

Synthetic polypeptide IT9302 was purchased from Kem-En-Tec A/S, Copenhagen, Denmark, with higher purity than 95%. This polypeptide was also conjugated to Keyhole Limpet Hemocyanin (KLH) by the producer. The soluble IT9302/KLH is more immunogenic than the polypeptide alone, and it may also be used as control protein for ELISA or Western Blotting.



*Immunization of rabbits*

250  $\mu$ g of IT9302 coupled to KLH, and as an emulsion with complete Freund's adjuvant, is used for intradermal or subcutaneous injections. The injections are repeated 4 times with  
5 two-week intervals and the rabbits are bled 8 and 12 days thereafter. Later booster injections of 250  $\mu$ g of IT9302/KLH in incomplete Freund's adjuvant are administered with one-month intervals using subcutaneous injections. The formation of IT9302 antibody is tested by dot-blot immunoassay or  
10 Western Blotting.

## REFERENCES

1. Bendtzen K. Lymphokines in inflammation. Inflammation Basic Mechanisms Tissue Injuring Principles and Clinical Models (P Venge & A Lindbom eds) 1985; Almqvist & Wiksell International. Stockholm: 187-217.
2. Bendtzen K. Interleukin-1, Interleukin-6, and tumor necrosis factor in infection, inflammation and immunity. Immunol Lett 1988;19:183-192.
3. Larsen C.G. Leukocyte activating and chemotactic cytokines in cell-mediated immune reactions of the human skin. Acta Dermatovenerol. 1991; Suppl. 160:1-48
4. Fiorentino D. F., M. W. Bond, and T. R. Mosmann. 1989. Two types of mouse helper T cell. IV. Th2 clones secrete a factor that inhibits cytokine production by Th1 clones. J. Exp. Med., 170:2081.
5. Viera P., R. de Wall-Malefyt, M.-N. Dang, K. E. Johnson, R. Kastelein, D. F. Fiorentino, J. E. de Vries, M.-G. Roncarolo, T. R. Mosmann, and K. W. Moore. 1991. Isolation and expression of human cytokine synthesis inhibitory factor (CSIF/IL-10) cDNA clones: homology to Epstein-Barr virus open reading frame BCRF1. Proc. Natl. Acad. Sci. (USA), 88:1172.
6. Moore, K.W., O'Garra A., de Waal Malefyt R., Vieira, Mosmann T.R. 1993. Interleukin-10, Annu Rev. Immunol, 11:165-90.
7. Kim, J.M., Brannan, C.I. Copeland N.G., Jenkins, N.A., Khan, T.A., Moore, K.W. 1992. Structure of the mouse interleukin-10 gene and chromosomal localization of the mouse and human genes. J. Immunol 148:3618-23.
8. Carter, D.B., Deibel, M.R-Jr, Dunn, C.J. et al. 1990. Purification, cloning, expression and biological characterization of an interleukin-1 receptor antagonist protein. NATURE 344:633-638.
9. Hannum, C.H., Wilcox, C.J., Arend, W.P. et al. 1990. Interleukin-1 receptor antagonist activity of a human interleukin-1 inhibitor. Nature 343:336-40.

10. Firestein, G.S., Boyle, D.L., Yu, C., et al. 1994. Synovial interleukin-1 receptor antagonist and interleukin-1 balance in rheumatoid arthritis. *Arthritis Rheum* 37:644-652.
- 5 11. Fisher, C.J.-Jr., Slotman, G.J., Opal, S.M., Pribble, J.P. et al. 1994. Initial evaluation of recombinant interleukin-1 receptor antagonist in the treatment of sepsis syndrome: a randomized, open-label, placebo-controlled multicenter trial. The IL-1RA Sepsis Syndrome Study Group. *Crit-Care-Med.* 22:12-21.
- 10 12. de Waal-Malefyt, R., Haanen J., Spits, H., et al. 1991. IL-10 and viral IL-10 strongly reduce antigen-specific human T cell proliferation by diminishing the antigen-presenting capacity of monocytes via down-regulation of class II MHC expression. *J. Exp. Med.* 174:915-24.
- 15 13. Gazzinelli, R.T., Oswald, I.P., James, S.L., Sher, A., 1992. IL-10 inhibits parasite killing and nitric oxide production by IFN- $\gamma$ -activated macrophages. *J. Immunol.* 148:1792-96.
- 20 14. Jinquan, T., Larsen, C.G., Gesser, B., Matsushima, K., Thestrup-Pedersen, K. 1993. Human IL-10 is a chemo-attractant for CD8+ T lymphocytes and an inhibitor of IL-8- induced CD4+ T lymphocyte Migration. *Journal of Immunology*, 151:4545-4551.
- 25 15. Rousset F., E. Garcia, T. Defrance, C. Peronne, D.-H. Hsu, R. Kastelein, K. W. Moore, and J. Banchereau. 1992. IL-10 is a potent growth and differentiation factor for activated human B lymphocytes. *Proc. Natl. Acad. Sci. USA*, 175:671.
- 30 16. Howard, M., O'Garra, A., Ishida, H., de Waal Malefyt, R., de Vries, J. 1992. Biological properties of Interleukin-10. *J. Clin. Immunol* 12:239-47.
17. Kuhn, R., Lohler, J., Rennick, D., Rajewsky, K., Muller, W. 1993. Interleukin-10-deficient mice develop chronic enterocolitis. *Cell* 75: 263-74.
- 35 18. Sher, A., Fiorentino, D.F., Caspar, P., Pearce, E., Mosmann, T. 1991. Production of IL-10 by CD4+ lymphocytes correlates with down-regulation of Th1 cytokine

- synthesis in helminth infection. *J. Immunol.* 147:2713-16.
19. Clerici, M., Shearer, G.M. 1993 *Immunology Today*. 14:107-111.
- 5 20. Bry, K., Lappalainen, U. 1994. Interleukin-4 and transforming growth factor-beta 1 modulate the production of interleukin-1 receptor antagonist and prostaglandin E2 by decidual cells. *Am-J-obstet-Gynecol* 170 (4): 1194-1198
- 10 21. Firestein, G., S., Boyle, D. L., Yu, C., Paine, M. M., Whisenand, T. D., Zvaifler, N. J., Arend, W. P. 1994. Synovial interleukin-1 receptor antagonist and interleukin-1 balance in rheumatoid arthritis. *Arthritis-Rheum*, 37/5: 644-652
- 15 22. Roberge, C. J., De-Medicis, R., Dayer, J. M., Rola-Pleszczynski, M., Naccahe, P. H., Poubelle, P. E. 1994. Crystal-induced neutrophil activation: V. Differential production of biologically active IL-1 receptor antagonist. *J. Immunol* 152/11: 5485-5494
- 20 23. McCall, R. D., Haskill, S., Zimmermann, E. M., Lund, P. K., Thompson, R. C., Sartor, R. B. 1994. Tissue interleukin 1 and interleukin-1 receptor antagonist expression in enterocolitis in resistant and susceptible rats. *Gastroenterology* (4): 960-72
- 25 25. Kimble, R. B., Vannice, J. L., Bloedow, D. C., Thompson, R. C., Hopfer, W., Kung, V. T., Brownfield, C., Pacifici, R. 1994. Interleukin-1 receptor antagonist decreases bone loss and bone resorption in ovariectomized rats. *J. Clin Invest.* 93/5: 1959-1967
- 30 26. Kline, J. N., Geist, L. J., Monick, M. M., Stinski, M. F., Hunninghake, G. W., 1994. *J. Immunol.* 152 (5): 2351-7
- 35 27. Tompkins, R. G. 1994. Human recombinant interleukin-1 receptor antagonist in the treatment of sepsis syndrome (editorial; comment). *Crit-Care-Med.* 22 (1): 3, 22 (1):12-21

28. Everaerd, B., Brouckaert, P., Fiers, W. 1994. Recombination IL-1 receptor antagonist protects against TNF-induced lethality in mice. *J. Immunol.* 152/10: 5041-5049
29. Fischer, C. J. Jr., Slotman, G. J., Opal, S. M.,  
5 Pribble, J. P., Bone, R. C., Emmanuel, G., Ng, D.,  
Bloedow, D. C., Catalano, M. A. 1994. Initial evaluation of human recombination interleukin-1 receptor antagonist in the treatment of sepsis syndrome: a randomized, open-label, placebo-controlled multicenter trial. The IL-1RA Sepsis Syndrome Study Group (see comments). *Crit-Care-Med.* 22(1): 12-21, 22(1): 3
- 10 30. Gomez-Reino-Carnoto, J. J. 1994. New therapies in rheumatoid arthritis. *Med-Clin* 543-545.
32. Nishihara, T., Ohsaki, Y., Ueda, N., Saito, N., Mundy,  
15 G. R. 1994. Mouse interleukin-1 receptor antagonist induced by actinobacillus actinomycetemcomitans lipopolysaccharide blocks the effects of interleukin-1 on bone resorption and osteoclast-like cell formation. *Infect-Immun.* 62(2): 390-7
- 20 33. Simon, C., Frances, A., Piquette, G. N., el-Danasouri, I., Zurawski, G., Dang, W., Polan, M. L. 1994. Embryonic implantation in mice is blocked by interleukin-1 receptor antagonist (see comments). *Endocrinology.* 134(2): 521-8, 134(2): 519-20
- 25 34. Baergen, R., Benirschke, K., Ulich, T. R., 1994. Cytokine expression in the placenta. The role of interleukin 1 and interleukin 1 receptor antagonist expression in chorioamnionitis and parturition. *Arch-Pathol-Lab-Med.* 118(1): 52-5
- 30 35. Tang, W.W., Feng, L., Vannice, J. L., Wilson, C. B. 1994. Interleukin-1 receptor antagonist ameliorates experimental antiglomerular basement membrane antibody-associated glomerulonephritis. *J. Clin-Invest.* 93(1): 279-9.
- 35 36. Cassatella, M. A., Meda, L., Gasperini, S., Calzetti, F., Bonara, S. 1994.
37. Interleukin 10 (IL-10) upregulates IL-1 receptor antagonist production from lipopolysaccharide-stimulated human

- polymorphonuclear leukocytes by delaying mRNA degradation. *J. Exp-Med.* 179/5: 1695-1699
38. Mancini, R., Bendetti, A., Jezequel, A. M. 1994. An interleukin-1 receptor antagonist decreases fibrosis induced by dimethylnitrosamine in rat liver. *Virchows-Arch.* 424/1: 25-31
39. Lukacs, N. W., Kunkel, S. L., Burdick, M. D., Lincoln, P. M., Strieter, R. M. 1993.
40. Interleukin-1 receptor antagonist blocks chemokine production in the mixed lymphocyte reaction. *Blood.* 82(12): 3668-74
41. Bandara, G., Mueller, G. M., Galea-Lauri, J., Tindal, M. H., Georgescu, H. I., Suchanek, M. K., Hung, G. L., Gloriso, J. C., Robbins, P. D., Evans, C. H. 1993.
42. Intraarticular expression of biologically active interleukin 1-receptor-antagonist protein by ex vivo transfer. *Proc-Natl-Acad-Sci-U-S-A.* 90(22): 10764-8
43. Dinarello, C. A. 1994. Anti-interleukin-1 strategies in the treatment of the septic shock syndrome. *Can-J-infect-Dis.* 5(suppl. A): 9A-16A
44. Oelmann, E., Topp, M. S., Reufi, B., Papadimitriou, C., Koeningsmann, M., Oberberg, D., Thiel, E., Berdel, W. E. 1994. *Int-J-Oncol.* 4/3: 555-558
45. Estrov, Z. 1993. Interruption of autocrine and paracrine growth-stimulatory mechanisms: a new therapeutic strategy for chronic myelogenous leukemia. *Semin-Hematol.* 30(3 suppl 3): 35-6
46. Wooley, P.H., Whalen, J.D., Chapman, D.L., Berger, A.E., Richard, K.A., Aspar, D.G., Staite, N.D. 1993. The effect of an interleukin-1 receptor antagonist protein on type II collagen-induced arthritis and antigen-induced arthritis in mice. *Arthritis Rheum.* 36 (9): 1305-1314
47. Peterson, C.M., Hales, H.A., Hatasaka, H.H., Mitchell, M.D., Rittenhouse, L., Jones, K.P. 1993. Interleukin-1 beta (IL-1 beta) modulates prostaglandin production and the natural IL-1 receptor antagonist inhibits ovulation

in the optimally stimulated rat ovarian perfusion model.  
Endocrinology 133 (5): 2301-2306

48. Estrov, Z., Kurzrock, R., Talpaz, M. 1993. Role of  
interleukin-1 inhibitory molecules in therapy of acute  
and chronic myelogenous leukemia. Leuk. Lymphoma 10 (6):  
407-418
49. Chensue, S.W., Bienkowski, M., Eessalu, T.E., Warming-  
ton, K.S., Hershey, S.D., Lukacs, N.W., Kunkel, S.L.  
1993. IL-1 receptor antagonist protein (IRAP)  
regulates schistosome egg granuloma formation and the  
regional lymphoid response. J. Immunol. 151 (7): 3654-  
3662
50. Bowyer, J.F., Davies, D.L., Schmued, L., Broening, H.W.,  
Newport, G.D., Slikker, W Jr., Holson, R.R. 1994. Fur-  
ther studies of the role of hyperthermia in methampheta-  
mine neurotoxicity. J. Pharmacol. Exp. Ther. 268/3:  
1571-1580
51. Cole, O.F., Sullivan, M.H.F., Elder, M.G. 1993. The  
'interleukin-1 receptor antagonist' is a partial agonist  
of prostaglandin synthesis by human decidual cells.  
Prostaglandins 46/6: 493-498
52. Jenkins, J.K., Arend, W.P. 1993. Interleukin 1 receptor  
antagonist production in human monocytes is induced by  
IL-1 $\alpha$ , IL-3, and IL-4 and GM-CSF. Cytokine 5/5: 407-  
415
53. Cocceani, F., Lees, J., Redford, J., Bishai, I. 1992.  
Interleukin-1 receptor antagonist: effectiveness against  
interleukin-1 fever. Can. J. Pharmacol. 70 (12): 1590-  
1596
54. Schiro, R., Longoni, D., Rossi, V., Maglia, O., Doni,  
A., Arsura, M., Carrara, G., Masera, G., Vannier, E.,  
Dinarello, C.A., Rambaldi, A., Biondi, A. 1994. Suppres-  
sion of juvenile chronic myelogenous leukemia colony  
growth by interleukin-1 receptor antagonist. Blood 83/2:  
460-465
55. Watson, M.L., Smith, D., Bourne, A.D., Thompson, R.C.,  
Westwick, J. 1993. Cytokines contribute to airway dys-  
function hyperreactivity, pulmonary eosinophil accumula-

- tion and tumor necrosis factor generation by pre-treatment with an interleukin-1 receptor antagonist. *Am. J. Respir. Cell Mol. Biol.* 8 (4): 365-369
56. Abhyankar, S., Gilliland, D.G., Ferrara, J.L.M. 1993. Interleukin-1 is a critical effector molecule during cytokine dysregulation in graft-versus-host disease to minor histocompatibility antigens. *Transplantation* 56/6: 1518-1523
57. Lan, H.Y., Nikolic Paterson, D.J., Zarama, M., Vannice, J.L., Atkins, R.C. 1993. Suppression of experimental crescentic glomerulonephritis by the interleukin-1 receptor antagonist. *Kidney Int.* 43 (2): 479-485
58. Herve, P. 1993. Prevention and treatment of acute GvHD - New modalities. *Nouv. Rev. Fr. Hematol.* 35/3: 295-297
59. Conti, P., Panara, M.R., Barbacane, R.C., Placido, F.C., Bongrazio, M., Reale, M., Dempsey, R.A., Fiore, S. 1992. Blocking the interleukin-1 receptor inhibits leukotriene B4 and prostaglandin E2 generation in human monocyte cultures. *Cell Immunol.* 145 (1): 199-209
60. Kristensen, M., Deleuran, B., Eedy, D.J., Feldmann, M., Breathnach, S.M., Brennan, F.M. 1992. Distribution of interleukin-1 receptor antagonist protein (IRAP), interleukin-1 receptor, and interleukin-1 alpha in normal and psoriatic skin, Decreased expression of IRAP in psoriatic lesional epidermis. *Br. J. Dermatol.* 127 (4): 305-311
- 6.1 Romero, R., Sepulveda, W., Mazor, M., Brandt, F., Cotton, D.B., Dinarello, C.A., Mitchell, M.D. 1992. The natural interleukin-1 receptor antagonist in term and pre-term parturition. *Am. J. Obstet. Gynecol.* 167 (4 Pt 1): 863-872
62. Dinarello, C.A. 1992. Reduction of inflammation by decreasing production of interleukin-1 or by specific receptor antagonism. *Int. J. Tissue. React.* 14 (2): 65-75
63. Conti, P., Panara, M.R., Barbacane, R.C., Bongrazio, M., Dempsey, R.A., Reale, M. 1993. Human recombinant IL-1 receptor antagonist (IL-1Ra) inhibits leukotriene B4



- generation from human monocyte suspensions stimulated by lipopolysaccharide (LPS). Clin. Exp. Immunol. 91/3: 526-531
64. DeForge, L.E., Tracey, D.E., Kenney, J.S., Remick, D.G.  
5 1992. Interleukin-1 receptor antagonist protein inhibits interleukin-8 expression in lipopolysaccharide-stimulated human whole blood. Am. J. Pathol. 140 (5): 1045-1054
65. Porat, R., Poutsika, D.D., Miller, L.C., Granowitz, E.V., Dinarello, C.A. 1992. Interleukin-1 (IL-1) receptor blockade reduces endotoxin and Borreliaburgdorferi-stimulated IL-8 synthesis in human monoclear cells. Faseb. J. 6 (7): 2482-2486
66. Boermeester, M.A., van Leeuwen, P.A.M., Schneider, A.J., Houdijk, A.P.J., Ferwerda, C.C., Wesdorp, R.I.C. 1993.  
15 Interleukin-1 receptor antagonist: A new therapeutic agent in the treatment of septic syndrome. Ned. Tijdschr. Geneesks. 137/7: 337-342
67. Smith, R.J., Chin, J.E., Sam, L.M., Justen, J.M. 1991. Biologic effects of an interleukin-1 receptor antagonist protein on interleukin-1-stimulated cartilage erosion and chondrocyte responsiveness. Arthritis Rheum. 34 (1):  
20 78-83
68. Conti, P., Barbacane, R.C., Panara, M.R., Reale, M., Placido, F.C., Fridas, S., Bongrazio, M., Dempsey, R.A.  
25 1992. Human recombinant interleukin-1 receptor antagonist (hrIL-1ra) enhances the stimulatory effect of interleukin-2 on natural killer cell activity against MOLT-4 target cells. Int. J. Immunopharm. 14/6: 987-993
69. Selig, W., Tocker, J. 1992. Effect of interleukin-1  
30 receptor antagonist on antigen-induced pulmonary responses in guinea pigs. Eur. J. Pharmacol. 213/3: 331-336
70. McCarthy, P.L. Jr., Abhyankar, S., Neben, S., Newman, G., Sieff, C., Thompson, R.C., Burakoff, S.J., Ferrara, J.L.M. 1991. Inhibition of interleukin-1 by an interleukin-1 receptor antagonist prevents graft-versus-host  
35 diseases. Blood 78/8: 1915-1918
71. Estrov, Z, Kurzrock, R., Wetzler, M., Kantarjian, H., Blake, M., Harris, D., Gutterman, J.U., Talpaz, M. 1991.

Suppression of chronic myelogenous leukemia colony growth by interleukin-1 (IL-1) receptor antagonist and soluble IL-1 receptors: A novel application for inhibitors of IL-1 activity. *Blood* 78/6: 1476-1484

- 5 72. Thomas, T.K., Will, P.C., Srivastava, A., Wilson, C.L., Harbison, M., Little, J., Chesonis, R.S., Pignatello, M., Schmolze, D., Symington, J., Kilin, P.L., Thompson, R.C. 1991. Evaluation of an interleukin-1 receptor antagonist in the rat acetic acid-induced colitis model.
- 10 Agents Actions 34/1-2: 187-190
73. Carter, D.B., Deibel, M.R. Jr., Dunn, C.J., Tomich, C.S.C., Laborde, A.L., Slightom, J.L., Berger, A.E., Bienkowski, M.J., Sun, F.F., McEwan, R.N., Harris, P.K.W., Yem, A.W., Waszak, G.A., Chosay, J.G., Sieu, L.C., Hardee, M.M., Zurcher Neely, H.A., Reardon, I.M.,
- 15 Heinrikson, R.L. et al. 1990. Purification, cloning expression and biological characterization of an interleukin-1 receptor antagonist protein. *Nature* 344/6267: 633-638
- 20 74. Larsen C.G., Anderson A.O., Apella E., Oppenheim J.J., Matsushima K., 1989. *Science* 243:1464;
75. Larsen C.G., Jinquan T., Deleurant B., Thestrup-Pedersen K. 1993, IL-10 is a potent regulator of the chemotactic response of mononuclear cells, but not of granulocytes. *J. Invest. Dermatol.* Vol 100, No 6
- 25 76. Sankoff and Kruskal in chapter 1 of "Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison" (Addison-Wesley, Reading, Mass. 1983).
- 30 77. Berzofsky, *Science* 229, (1985) 932-940
78. Bowie et al., *Science* 247, (1990) 1306-1310
79. Wasserman et al., *J. Immunol.* 87, 1961, 290-295
80. Levine et al., *Methods in Enzymology* 11, 1967, 928-936
81. Lewis et al., *Biochemistry* 22, 1983, 948-954
- 35 82. Rene de Waal Malefyt, John Abrahams, Bruce Bennet, Carl G. Figdor and Jan E. de Vries (1991), Interleukin 10 (IL-10) Inhibits Cytokine Synthesis by Human Monocytes:

- An Autoregulatory Role of IL-10 Produced by Monocytes.  
*J. Exp. Med.* 174, 1209-1220
83. Szoka et al., *Ann. Rev. Biophys. Bioeng.* 9, 1980, 467
84. US Patent No. 4,235,871
- 5 85. US 4,501,728
86. US 4,837,028.
87. Walter H. Gotlieb, John S. Abrams, Joanna M. Watson,  
Thierry J. Velu, Jonathan S. Berek, Otoniel Martinez-  
Meza (1992), Presence of interleukin 10 (IL-10) in the  
10 ascites of patients with ovarian and other intra-abdomi-  
nal cancers. *Cytokine* 4, No. 5, 385-390
88. Blanco G., P. Gianello, Sh. Germana, M. Baetscher, D.H.  
Sachs and Chr. LeGuern (1995), Molecular identification  
of porcine interleukin 10: Regulation of expression in  
15 kidney allograft model. *Proc. Natl. Acad. Sci. USA* 92,  
2800-2804

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Nycomed DAK A/S
- (B) STREET: P.O. Box 1911, Lergraversvej 59
- (C) CITY: Copenhagen S
- (E) COUNTRY: Denmark
- (F) POSTAL CODE (ZIP): 2300

(ii) TITLE OF INVENTION: Immunomodulators

(iii) NUMBER OF SEQUENCES: 23

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Ala Tyr Met Thr Met Lys Ile Arg Asn  
1 5

## (2) INFORMATION FOR SEQ ID NO: 2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Ala Tyr Met Thr Ile Lys Met Arg Asn  
1 5

(2) INFORMATION FOR SEQ ID NO: 3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

Ala Phe Met Thr Leu Lys Leu Arg Asn  
1 5

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Ala Tyr Met Thr Met Lys Val Arg Glu  
1 5

(2) INFORMATION FOR SEQ ID NO: 5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

Gly Tyr Met Thr Met Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

Ala Phe Met Thr Met Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

Ala Tyr Ile Thr Met Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Ala Tyr Leu Thr Met Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Ala Tyr Val Thr Met Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Ala Tyr Met Thr Ile Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

Ala Tyr Met Thr Leu Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 9 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

Ala Tyr Met Thr Val Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 13:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 13:

Ala Tyr Met Thr Met Lys Ile Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 14:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 14:

Ala Tyr Met Thr Met Lys Met Arg Asp  
1 5

(2) INFORMATION FOR SEQ ID NO: 15:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS:  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 15:

Ala Tyr Met Thr Met Lys Val Arg Asp  
1 5



## (2) INFORMATION FOR SEQ ID NO: 16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide

- (iii) HYPOTHETICAL: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 16:

Ala Tyr Met Thr Met Lys Ile Asp Gln  
1 5

## (2) INFORMATION FOR SEQ ID NO: 17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide

- (iii) HYPOTHETICAL: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 17:

Ala Tyr Met Thr Met Lys Ile Asp Glu  
1 5

## (2) INFORMATION FOR SEQ ID NO: 18:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: peptide

- (iii) HYPOTHETICAL: NO

- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 18:

Ser Pro Gly Gln Gly Thr Gln Ser Glu  
1 5

## (2) INFORMATION FOR SEQ ID NO: 19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 6 amino acids

- (B) TYPE: amino acid
- (C) STRANDEDNESS:
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 19:

Thr Xaa Lys Xaa Arg Xaa  
1 5

(2) INFORMATION FOR SEQ ID NO: 20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 7 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 20:

Xaa Thr Xaa Lys Xaa Arg Xaa  
1 5

(2) INFORMATION FOR SEQ ID NO: 21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 8 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 21:

Xaa Xaa Thr Xaa Lys Xaa Arg Xaa  
1 5

(2) INFORMATION FOR SEQ ID NO: 22:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS:
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 22:

Xaa Xaa Xaa Thr Xaa Lys Xaa Arg Xaa  
1 5

(2) INFORMATION FOR SEQ ID NO: 23:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 23:

GCCTACATGA CAATGAAGAT ACGAAAC

## CLAIMS

1. A polypeptide other than human interleukin 10 which has at least one of the following properties:

- 5 a) induces inhibition of spontaneous IL-8 production by human monocytes,
- b) induces inhibition of IL-1 $\beta$  induced IL-8 production by human peripheral blood mononuclear cells (PBMC),
- c) induces production of interleukin-1 receptor antagonistic protein (IRAP) by human monocytes,
- 10 d) induces chemotactic migration of CD8+ human T lymphocytes *in vitro*,
- e) desensitizes human CD8+ T cells resulting in an unresponsiveness towards rhIL-10,
- f) suppresses the chemotactic response of CD4+ T human lymphocytes towards IL-8,
- 15 g) suppresses the chemotactic response of human monocytes towards MCAF/MCP-1,
- h) does not inhibit class II MHC molecule expression on human monocytes, in contrast to human IL-10.
- 20 i) induces the production of IL-4 by cultured normal human CD4+ T cells,
- j) reduces the TNF $\alpha$  production in human mixed leukocyte reaction.

2. A polypeptide having the formula

25 Thr-X<sub>4</sub>-Lys-X<sub>5</sub>-Arg-X<sub>6</sub> (SEQ ID NO:19),

wherein

X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and

30 X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.

3. A polypeptide having the formula


X<sub>3</sub>-Thr-X<sub>4</sub>-Lys-X<sub>5</sub>-Arg-X<sub>6</sub> (SEQ ID NO:20),

wherein

X<sub>3</sub>, X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and

5 X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.

4. A polypeptide having the formula




wherein

X<sub>2</sub> is Tyr or Phe,

10 X<sub>3</sub>, X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and

X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.

5. A polypeptide having the formula



15 wherein

X<sub>1</sub> is Ala or Gly,

X<sub>2</sub> is Tyr or Phe,

X<sub>3</sub>, X<sub>4</sub> and X<sub>5</sub> are independently selected from the group consisting of Met, Ile, Leu and Val; and

20 X<sub>6</sub> is selected from the group consisting of Asp, Gln and Glu.

6. A polypeptide according to any of claims 2-5, wherein X<sub>1</sub> is Ala.

7. A polypeptide according to any of claims 2-5, wherein X<sub>1</sub> is Gly.

25 8. A polypeptide according to any of claims 2-7, wherein X<sub>2</sub> is Tyr.

9. A polypeptide according to any of claims 2-7, wherein  $X_2$  is Phe.
10. A polypeptide according to any of claims 2-9, wherein  $X_3$  is Met.
- 5 11. A polypeptide according to any of claims 2-9, wherein  $X_3$  is Ile.
12. A polypeptide according to any of claims 2-9, wherein  $X_3$  is Leu.
- 10 13. A polypeptide according to any of claims 2-9, wherein  $X_3$  is Val.
14. A polypeptide according to any of claims 2-13, wherein  $X_4$  is Met.
15. A polypeptide according to any of claims 2-13, wherein  $X_4$  is Ile.
- 15 16. A polypeptide according to any of claims 2-13, wherein  $X_4$  is Leu.
17. A polypeptide according to any of claims 2-13, wherein  $X_4$  is Val.
- 20 18. A polypeptide according to any of claims 2-17, wherein  $X_5$  is Met.
19. A polypeptide according to any of claims 2-17, wherein  $X_5$  is Ile.
20. A polypeptide according to any of claims 2-17, wherein  $X_5$  is Leu.
- 25 21. A polypeptide according to any of claims 2-17, wherein  $X_5$  is Val.

22. A polypeptide according to any of claims 2-21, wherein X<sub>6</sub> is Asp.

23. A polypeptide according to any of claims 2-21, wherein X<sub>6</sub> is Gln.

5 24. A polypeptide according to any of claims 2-21, wherein X<sub>6</sub> is Glu.

25. A substance or polypeptide which comprises an amino acid sequence as defined in any of claims 2-24.

10 26. A polypeptide comprising a subsequence of the sequence of human IL-10 and which has a degree of homology to vIL-10 which is 66% and a degree of homology to mIL-10 which is 44%.

27. Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn.

28. A polypeptide selected from the group consisting of

Ala-Tyr-Met-Thr-Ile-Lys-Met-Arg-Asn (SEQ ID NO:2),  
15 Ala-Phe-Met-Thr-Leu-Lys-Leu-Arg-Asn (SEQ ID NO:3),  
Ala-Tyr-Met-Thr-Met-Lys-Val-Arg-Glu (SEQ ID NO:4),  
Gly-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:5),  
Ala-Phe-Met-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:6),  
Ala-Tyr-Ile-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:7),  
20 Ala-Tyr-Leu-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:8),  
Ala-Tyr-Val-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:9),  
Ala-Tyr-Met-Thr-Ile-Lys-Ile-Arg-Asp (SEQ ID NO:10),  
Ala-Tyr-Met-Thr-Leu-Lys-Ile-Arg-Asp (SEQ ID NO:11),  
Ala-Tyr-Met-Thr-Val-Lys-Ile-Arg-Asp (SEQ ID NO:12),  
25 Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:13),  
Ala-Tyr-Met-Thr-Met-Lys-Met-Arg-Asp (SEQ ID NO:14),  
Ala-Tyr-Met-Thr-Met-Lys-Val-Arg-Asp (SEQ ID NO:15),  
Ala-Tyr-Met-Thr-Met-Lys-Ile-Asp-Gln (SEQ ID NO:16), and  
Ala-Tyr-Met-Thr-Met-Lys-Ile-Asp-Glu (SEQ ID NO:17).

29. A substance or polypeptide according to any of claims 1-28 in substantially pure form.
30. A nucleotide sequence encoding a polypeptide according to any of claims 2-29.
- 5 31. An antibody which binds to a substance or polypeptide according to any of claims 1-29.
32. An antibody which binds specifically to a polypeptide having the amino acid sequence SEQ ID NO:1.
- 10 33. An antibody according to claim 31 or 32 which is a polyclonal antibody.
34. An antibody according to claim 31 or 32 which is a monoclonal antibody.
- 15 35. An antibody according to any of claims 31-34 which is capable of neutralizing one or more of the activities a) to g) mentioned in claim 1.
36. A substance other than the monoclonal antibody 19F1 which is capable of neutralizing one or more of the activities a) to j) mentioned in claim 1.
- 20 37. A pharmaceutical composition comprising a substance according to claim 36.
38. Use of a substance according to claim 36 for diminishing or neutralizing high concentration of hIL-10 and/or vIL-10.
- 25 39. Use of a substance according to claim 36 for the manufacture of a pharmaceutical composition for the treatment or prophylaxis of ovary cancer and/or AIDS.
40. A method of treating and/or preventing ovary cancer and/or AIDS, the method comprising administering to a patient



in need thereof a therapeutically or prophylactically effective amount of a substance according to claim 36.

41. Use of a compound which has an hIL-10 antagonist activity for the manufacture of a pharmaceutical composition for  
5 treatment or prophylaxis of ovary cancer and/or AIDS.

42. A method of treating and/or preventing ovary cancer and/or AIDS, the method comprising administering to a patient in need thereof a therapeutically or prophylactically effective amount of a compound which has a hIL-10 antagonist  
10 activity.

43. A pharmaceutical composition comprising a substance or polypeptide according to any of claims 1-29.

44. Use of a substance or polypeptide according to any of claims 1-29 for the treatment or prophylaxis of one or more  
15 of the diseases mentioned in Table 3.

45. Use of a substance or polypeptide according to any of claims 1-29 for the manufacture of a pharmaceutical composition for the treatment or prophylaxis of one or more of the diseases mentioned in Table 3.

20 46. A method of treating and/or preventing one or more of the diseases mentioned in Table 3, the method comprising administering to a patient in need thereof a therapeutically or prophylactically effective amount of a substance or polypeptide according to any of claims 1-29.

## AMENDED CLAIMS

[received by the International Bureau on 6 November 1995 (06.11.95);  
original claim 28 amended; remaining claims unchanged (1 page)]

22. A polypeptide according to any of claims 2-21, wherein  $X_6$  is Asp.
23. A polypeptide according to any of claims 2-21, wherein  $X_6$  is Gln.
- 5 24. A polypeptide according to any of claims 2-21, wherein  $X_6$  is Glu.
25. A substance or polypeptide which comprises an amino acid sequence as defined in any of claims 2-24.
26. A polypeptide comprising a subsequence of the sequence of  
10 human IL-10 and which has a degree of homology to vIL-10 which is 66% and a degree of homology to mIL-10 which is 44%.
27. Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asn.
28. A polypeptide selected from the group consisting of
- Ala-Tyr-Met-Thr-Ile-Lys-Met-Arg-Asn (SEQ ID NO:2),  
15 Ala-Phe-Met-Thr-Leu-Lys-Leu-Arg-Asn (SEQ ID NO:3),  
Ala-Tyr-Met-Thr-Met-Lys-Val-Arg-Glu (SEQ ID NO:4),  
Gly-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:5),  
Ala-Phe-Met-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:6),  
Ala-Tyr-Ile-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:7),  
20 Ala-Tyr-Leu-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:8),  
Ala-Tyr-Val-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:9),  
Ala-Tyr-Met-Thr-Ile-Lys-Ile-Arg-Asp (SEQ ID NO:10),  
Ala-Tyr-Met-Thr-Leu-Lys-Ile-Arg-Asp (SEQ ID NO:11),  
Ala-Tyr-Met-Thr-Val-Lys-Ile-Arg-Asp (SEQ ID NO:12),  
25 Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Asp (SEQ ID NO:13),  
Ala-Tyr-Met-Thr-Met-Lys-Met-Arg-Asp (SEQ ID NO:14),  
Ala-Tyr-Met-Thr-Met-Lys-Val-Arg-Asp (SEQ ID NO:15),  
Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Gln (SEQ ID NO:16), and  
Ala-Tyr-Met-Thr-Met-Lys-Ile-Arg-Glu (SEQ ID NO:17).

1/13




FIG. 1. Comparison of the predicted amino acid sequences of mIL-10, hIL-10, and BCRF1. Amino acid sequence identities are indicated by vertical lines.

**Fig. 1**

2/13

COOH TERMINAL PEPTIDE SEQUENCES OF IL-10 INCLUDING  
9 AMINO ACIDS OF PORCINE, HUMAN and VIRUS PROTEINS.

PORCINE IL-10				*		*		*												
PIT	-	A	-	Y	-	M	-	T	-	M	-	K	-	M	-	R	-	K	-	N
HUMAN IL-10				*		*		*												
IT 9302	-	A	-	Y	-	M	-	T	-	M	-	K	-	I	-	R	-	N		
BCRFI IL-10				*		*		*												
VIT	-	A	-	Y	-	M	-	T	-	I	-	K	-	A	-	R	-			


Aminoacids marked with \* represent key position for  
receptor binding and are evolutionary well conserved.  
Compare the IL-10 aminoacid sequence from mouse below.

MOUSE IL-10						*														
MIT	-	A	-	Y	-	M	-	M	-	I	-	K	-	M	-	K	-	S		

This sequence is lacking the homology to the important  
amino acids.

Fig. 2

3/13



IT9302 inhibits spontaneous IL-8 production by purified cultured monocytes. (▲) Indicates the level of IL-8 when using rh IL-10 (100 ng/ml).

**Fig. 3**

4/13




Fig. 4

5/13

IT9302 stimulated human monocytes




Fig. 5

6/13





Fig. 6



7/13

Chemotactic activity of IT9302  
on CD8+ T cells




IT9302 (ng/ml)

Fig. 7

8/13


Desensitization of CD8+ T cells by  
IT9302, resulting in unresponsiveness  
of CD8+ T cells towards IL-10 (10 ng/ml) induced chemotaxis

**Fig. 8**

9/13

# Suppression of IL-8 activity by IT9302


Effect of IT9302 on IL-8 (10 ng/ml) mediated chemotaxis.



**Fig. 9**

10/13


IT9302 inhibits MCAF/MCP-1  
induced monocyte chemotaxis



**Fig. 10**

11/13

ECL-Western Blotting of CD4+ T cell cytosolic proteins using a goat anti-human IL-4 antibody.




Stimulation for 3 days with:

1. Control
2. mon.anti-IL-8 antibody WS.4 10  $\mu$ g/ml,
3. rIL-8 100 ng/ml,
4. rIL-10 100 ng/ml,
5. IT9302 10 ng/ml,
6. rIFN gamma 10 ng/ml.

Fig. 11

12/13

ECL-Western Blotting of Human Mixed lymphocyte culture cytosolic proteins using a rabbit anti-human TNF- $\alpha$  antibody.



Stimulation for 24 hours with:

1. Control
2. IT9302 10 ng/ml

Fig. 12

13/13




Fig. 13

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/DK 95/00227

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/24 C07K14/54 C07K7/06 C07K16/34 A61K38/20  
A61K39/395

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X  A	<p>WO,A,91 00349 (SCHERING CORPORATION) 10 January 1991</p> <p>see page 3, line 15 - page 5, line 2 see page 13, line 26 - page 14, line 30 see page 16, line 20 - page 17, line 20 see page 21, line 30 - line 36; claims; example VI &amp; EP,A,0 405 980 cited in the application</p> <p style="text-align: center;">---</p> <p style="text-align: center;">-/--</p>	<p>25, 29-31, 33,34, 43-46 1-24, 26-28, 32,35-42</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

22 August 1995

Date of mailing of the international search report

29.08.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Montero Lopez, B



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/DK 95/00227

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X  A	<p>WO,A,93 02693 (SCHERING CORPORATION) 18 February 1993 cited in the application</p> <p>see page 4, line 25 - page 7, line 10 see page 21, line 3 - page 24, line 9 see page 32, line 6 - line 25; examples 1-3</p> <p style="text-align: center;">---</p>	<p>25, 29-31, 33,34, 43-46 1-24, 26-28, 32,35-42</p>
X	<p>WO,A,93 17698 (SCHERING CORPORATION) 16 September 1993 see page 3, line 2 - line 12 see page 6, line 23 - page 7, line 9 see SEQ ID Nos. 1-4</p> <p style="text-align: center;">-----</p>	<p>25,29, 30,43-46</p>

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK95/00227

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 40, 42, 44, 46  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 40, 42, 44 and 46 are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/DK 95/00227

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9100349	10-01-91	AU-B- 635058	11-03-93
		AU-A- 6077090	17-01-91
		CA-A- 2062763	29-12-90
		CN-A- 1051393	15-05-91
		EP-A- 0405980	02-01-91
		EP-A- 0567450	03-11-93
		JP-T- 4502560	14-05-92
		US-A- 5231012	27-07-93
WO-A-9302693	18-02-93	AU-A- 2441192	02-03-93
		CA-A- 2115060	18-02-93
		CZ-A- 9400243	18-01-95
		EP-A- 0541214	12-05-93
		EP-A- 0600970	15-06-94
		FI-A- 940519	04-02-94
		HU-A- 66865	30-01-95
		JP-T- 7502019	02-03-95
		NO-A- 940370	04-02-94
WO-A-9317698	16-09-93	CA-A- 2131524	16-09-93
		CN-A- 1079166	08-12-93
		EP-A- 0629130	21-12-94
		JP-T- 7504437	18-05-95
		ZA-A- 9301489	28-10-93